

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US99/19443

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : C07H 21/02; C12N 15/82; A01H 5/00
US CL : 536/23.1; 800/281, 298; 435/69.1, 419, 468

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 536/23.1; 800/281, 298; 435/69.1, 419, 468

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)
WEST/DERWENT, GENBANK, EMBL

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 5,777,201 A (POUTRE et al) 07 July 1998, especially col.14-18.	1-5,9-11,13-15
Y		----- 6-8,12,16-21
Y	US 5,057,419 A (MARTIN et al) 15 October 1991, col. 7-11.	1-21
Y	STUKEY et al. The OLE1 Gene of Saccharomyces cerevisiae Encodes the delta 9 Fatty Acid Desaturase and Can Be Functionally Replaced by the Rat Stearoyl-CoA Desaturase. Gene. The Journal of Biological Chemistry. 25 November 1990, Vol. 265, No. 33, pages 20144-20149, especially page 20146.	1-21

☒ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
E earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*G* document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means	
P document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search 12 NOVEMBER 1999	Date of mailing of the international search report 06 DEC 1999
Name and mailing address of the ISA/US Commissioner of Patents and Trademarks Box PCT Washington, D.C. 20231 Facsimile No. (703) 305-3230	Authorized officer <i>D. Lawrence Fox</i> ELIZABETH F. MCELWAIN Telephone No. 308-0196

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US99/19443

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X - Y	POLASHOCK et al. Expression of the Yeast delta-9 Fatty Acid Desaturase in Nicotiana tabacum. Plant Physiology. 1992, Vol. 100, pages 894-901, especially page 895.	1-5,9-11,13-15 ----- 6-8,12,16-21

PCT

INTERNATIONAL PRELIMINARY EXAMINATION REPORT

(PCT Article 36 and Rule 70)


Applicant's or agent's file reference RUT 97-0081/	FOR FURTHER ACTION	See Notification of Transmittal of International Preliminary Examination Report (Form PCT/IPEA/416)
International application No. PCT/US99/19443	International filing date (day/month/year) 24 AUGUST 1999	Priority date (day/month/year) 24 AUGUST 1998
International Patent Classification (IPC) or national classification and IPC Please See Supplemental Sheet.		
Applicant RUTGERS, THE STATE UNIVERSITY OF NEW JERSEY		

1. This international preliminary examination report has been prepared by this International Preliminary Examining Authority and is transmitted to the applicant according to Article 36.
2. This REPORT consists of a total of 4 sheets.
- ☐ This report is also accompanied by ANNEXES, i.e., sheets of the description, claims and/or drawings which have been amended and are the basis for this report and/or sheets containing rectifications made before this Authority. (see Rule 70.16 and Section 607 of the Administrative Instructions under the PCT).

These annexes consist of a total of 0 sheets.

3. This report contains indications relating to the following items:

- I ☒ Basis of the report
- II ☐ Priority
- III ☐ Non-establishment of report with regard to novelty, inventive step or industrial applicability
- IV ☐ Lack of unity of invention
- V ☒ Reasoned statement under Article 35(2) with regard to novelty, inventive step or industrial applicability; citations and explanations supporting such statement
- VI ☐ Certain documents cited
- VII ☐ Certain defects in the international application
- VIII ☐ Certain observations on the international application

Date of submission of the demand 15 MARCH 2000	Date of completion of this report 28 DECEMBER 2000
Name and mailing address of the IPEA/US Commissioner of Patents and Trademarks Box PCT Washington, D.C. 20231 Facsimile No. (703) 305-3230	Authorized officer  ELIZABETH F. MCELWAIN Telephone No. 308-0196

INTERNATIONAL PRELIMINARY EXAMINATION REPORT

International application No.

PCT/US99/19443

I. Basis of the report

1. With regard to the elements of the international application: *

☒ the international application as originally filed☒ the description:

pages 1-62

pages NONE, as originally filed

pages NONE, filed with the demand

pages NONE, filed with the letter of

☒ the claims:

pages 63-66

pages NONE, as originally filed

pages NONE, as amended (together with any statement) under Article 19

pages NONE, filed with the demand

pages NONE, filed with the letter of

☒ the drawings:

pages 1-2

pages NONE, as originally filed

pages NONE, filed with the demand

pages NONE, filed with the letter of

☒ the sequence listing part of the description:

pages 1-5

pages NONE, as originally filed

pages NONE, filed with the demand

pages NONE, filed with the letter of

2. With regard to the **language**, all the elements marked above were available or furnished to this Authority in the language in which the international application was filed, unless otherwise indicated under this item.
These elements were available or furnished to this Authority in the following language _____ which is:☐ the language of a translation furnished for the purposes of international search (under Rule 23.1(b)).☐ the language of publication of the international application (under Rule 48.3(b)).☐ the language of the translation furnished for the purposes of international preliminary examination (under Rules 55.2 and/or 55.3).3. With regard to any **nucleotide and/or amino acid sequence** disclosed in the international application, the international preliminary examination was carried out on the basis of the sequence listing:☒ contained in the international application in printed form.☒ filed together with the international application in computer readable form.☐ furnished subsequently to this Authority in written form.☐ furnished subsequently to this Authority in computer readable form.☐ The statement that the subsequently furnished written sequence listing does not go beyond the disclosure in the international application as filed has been furnished.☐ The statement that the information recorded in computer readable form is identical to the written sequence listing has been furnished.4. ☒ The amendments have resulted in the cancellation of:☒ the description, pages NONE☒ the claims, Nos. NONE☒ the drawings, sheets/fig. NONE5. ☐ This report has been drawn as if (some of) the amendments had not been made, since they have been considered to go beyond the disclosure as filed, as indicated in the Supplemental Box (Rule 70.2(c)).**

* Replacement sheets which have been furnished to the receiving Office in response to an invitation under Article 14 are referred to in this report as "originally filed" and are not annexed to this report since they do not contain amendments (Rules 70.16 and 70.17).

**Any replacement sheet containing such amendments must be referred to under item 1 and annexed to this report.

INTERNATIONAL PRELIMINARY EXAMINATION REPORT

International application No.

PCT/US99/19443

V. Reasoned statement under Article 35(2) with regard to novelty, inventive step or industrial applicability; citations and explanations supporting such statement

1. statement

Novelty (N)	Claims <u>1-21</u>	YES
	Claims <u>NONE</u>	NO
Inventive Step (IS)	Claims <u>NONE</u>	YES
	Claims <u>1-21</u>	NO
Industrial Applicability (IA)	Claims <u>1-21</u>	YES
	Claims <u>NONE</u>	NO

2. citations and explanations (Rule 70.7)

Claims 1-21 lack an inventive step under PCT Article 33(3) as being obvious over Poutre et al or Polashock et al in view of Applicants' admitted state of the prior art.
Poutre et al and Polashock et al each teach transforming a plant with a yeast desaturase gene.
Neither Poutre et al or Polashock et al teach customizing the sequence for improved expression in a plant cytoplasm.
Applicants' admitted state of the prior art teaches that modifying sequences for improved expression in plants was known (see page 18, at least)
Given the recognition of those of ordinary skill in the art of the value of expressing a yeast desaturase in plant for the purpose of modifying fatty acid expression in the plant, it would have been obvious to one of ordinary skill in the art to use the methods of Poutre et al or Polashock et al and to modify these methods by substituting a gene that has been customized for expression in a plant cytoplasm.

----- NEW CITATIONS -----
NONE

INTERNATIONAL PRELIMINARY EXAMINATION REPORT

International application No.

PCT/US99/19443

Supplemental Box

(To be used when the space in any of the preceding boxes is not sufficient)

Sheet 10

Continuation of: Boxes I - VIII

CLASSIFICATION:

The International Patent Classification (IPC) and/or the National classification are as listed below:
IPC(7): C07H 21/02; C12N 15/82; A01H 5/00 and US Cl.: 536/23.1; 800/281, 298; 435/69.1, 419, 468

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶: C07H 21/02, C12N 15/82, A01H 5/00	A1	(11) International Publication Number: WO 00/11012 (43) International Publication Date: 2 March 2000 (02.03.00)
(21) International Application Number: PCT/US99/19443 (22) International Filing Date: 24 August 1999 (24.08.99) (30) Priority Data: 60/097,586 24 August 1998 (24.08.98) US (71) Applicant (for all designated States except US): RUTGERS, THE STATE UNIVERSITY OF NEW JERSEY [US/US]; Old Queens, Somerset Street, New Brunswick, NJ 08903 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): MARTIN, Charles, E. [US/US]; 3 Seely's Run, Somerset, NJ 08873 (US). MITCHELL, Andrew [GB/US]; 106 North 3rd Street, Highland Park, NJ 08904 (US). (74) Agents: REED, Janet, E. et al.; Dann, Dorfman, Herrell and Skillman, Suite 720, 1601 Market Street, Philadelphia, PA 19103-2307 (US).	(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i>	
(54) Title: SYNTHETIC FATTY ACID DESATURASE GENE FOR EXPRESSION IN PLANTS (57) Abstract A synthetic fatty acid desaturase gene for expression in a multicellular plant is provided. The gene comprises a desaturase domain and a cyt b ₅ domain, and is customized for expression in a plant cytoplasm. Methods for designing and making a synthetic fatty acid desaturase gene customized for expression in a plant cytoplasm are also provided.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

**SYNTHETIC FATTY ACID DESATURASE GENE
FOR EXPRESSION IN PLANTS**

This application claims priority to U.S. Provisional Application No. 60/097,586, filed August 24, 1998, the entirety of which is incorporated by reference herein.

5

FIELD OF THE INVENTION

This invention relates to the field of genetic engineering, and more particularly to transformation of plants with heterologous fatty acid desaturase genes modified for optimum expression in plants.

10

BACKGROUND OF THE INVENTION

Several publications are referenced in this application in order to more fully describe the state of the art to which this invention pertains. The disclosure of each of these publications is incorporated by reference herein.

15

Alteration of fatty acid desaturation in plants is of interest to plant biologists and food scientists alike, due to the influence of unsaturated fatty acids on the health benefits and flavors of foods, as well as the role of these molecules in plant biological processes. For a nation interested in healthy diet, the quality of fats and oils depends on their fatty acid composition, with oils high in monounsaturated fatty acids (e.g., canola, olive) gaining popularity as new health benefits are discovered. Considering the flavors of plant foods, many flavor-producing compounds are derived from peroxidation of unsaturated fatty acids. Thus, efforts are being made to produce plants with increased amounts

20

25

30

-2-

of unsaturated fatty acids, preferably monounsaturated fatty acids.

In animal and fungal cells, monounsaturated fatty acids are aerobically synthesized from saturated fatty acids by a microsomal Δ -9 fatty acid desaturase that is membrane bound and cytochrome b_5 -dependent. A double bond is inserted between the 9- and 10-carbons of palmitoyl (16:0) and stearoyl (18:0) CoA to form palmitoleic (16:1) and oleic (18:1) acids. In the reaction mechanism, electrons are transferred from NADH-dependent cytochrome b_5 reductase, via the heme-containing cytochrome b_5 (Cyt b_5) molecule, to the Δ -9 fatty acid desaturase. The major form of cytochrome b_5 in animal, fungal and plant cells exists as an independent protein molecule that is anchored to the membrane by a short, carboxyl terminal, hydrophobic stretch of amino acids. The carboxyl terminal anchor orients the heme group of the Cyt b_5 on the membrane surface and allows it to translationally diffuse across the surface of the membrane. This property of lateral mobility allows this form of cytochrome b_5 to participate as an electron donor to a number of different proteins that catalyze a variety of metabolic reactions on the membrane surface, including fatty acid desaturases, various sterol biosynthetic enzymes and a variety of cytochrome P450 mediated reactions. While this contributes to the versatility of Cyt b_5 as an electron donor, it also implies that the major form of cytochrome b_5 shuttles between its redox partners by translational diffusion across the surface of the membrane (Strittmatter and Rogers, Proc. Natl. Acad. Sci. USA, 72: 2658-2661, (1975; Lederer, Biochimie 76: 674-692, 1994). Furthermore, this mechanism suggests that an independent, membrane bound cytochrome b_5 molecule

-3-

can potentially limit the rate of the metabolic reaction, depending on its abundance, its location on the membrane surface, its proximity to the electron acceptor, and the rate at which it can move and orient itself to the acceptor on the membrane surface.

In plants, unsaturated fatty acids are formed and incorporated into complex lipids in two distinct cellular compartments. *De novo* fatty acid synthesis occurs almost exclusively in the plastids, producing the saturated species 16:0-ACP (acyl carrier protein) and 18:0-ACP. 18:1-ACP is formed from 18:0-ACP in the plastid by a soluble, ferredoxin-dependent Δ -9 desaturase. These fatty acids are then shunted into one of two routes - a plastid-localized "procaryotic" pathway or a cytosolic/ER (endoplasmic reticulum) "eucaryotic" pathway - for further modification and acylation into glycerolipids (Somerville and Browse, *Science* 252: 80-87, 1991). The acyl ACPs that are shunted into the prokaryotic pathway remain within the plastid and are used for the synthesis of phosphatidic acid and further conversion to chloroplast glycerolipids. The fatty acyl groups of those lipids may be further desaturated by plastid desaturases that also use ferredoxin as the electron donor.

Acyl-ACPs that are shunted into the eukaryotic pathway are converted to free fatty acids, transported across the chloroplast membrane into the cytoplasm where they are converted to acyl CoA thioesters by acyl CoA synthetase. Those fatty acids are then converted to cytoplasmic/ER phosphatidic acid which can then be converted to membrane glycerophospholipids, or storage lipids, in the form of triacylglycerols and sterol esters that are the major components of plant oils.

-4-

Most polyunsaturated 18-carbon plant fatty acids appear to be formed in the cytosol by the ER-bound desaturases (Table 1). Once the 18:1 fatty acid is incorporated into phospholipid, an ER-bound desaturase can catalyze the formation of a Δ -12 double bond in the fatty acyl chain to form Δ -9,12 18:2. Other ER bound desaturase enzymes can act on 18:2 to introduce a Δ -15 double bond to form Δ 9,12,15 18:3. These desaturase are thought to be similar to animal and fungal desaturases because they are membrane bound and appear to require a cytochrome b_5 -mediated electron transport chain.

TABLE 1:

Plant	Gene	Desaturase Type	Primary Activity	b5 chimera	Reference
Arabidopsis	FAD2	Δ 12, microsomal	18:1->18:2	no	Okuley J. et al. Plant Cell 6: 147-158, 1994
Arabidopsis	FAD3	Δ 15, microsomal	18:2->18:3	no	Shah S. & Z. Xin, Plant Physiol. 114: 1533-1539, 1997
Nicotiana tabacum	NtFAD3	Δ 15, microsomal	18:2->18:3	no	Hamada T. et al. Plant & Cell. Physiol. 37: 606-611, 1996, Hamada T. et al. Transgenic Res. 5: 115-121, 1996
Soybean	FAD 2-1	Δ 12, microsomal, developing seeds	18:1->18:2	no	Heppard E.P. et al. Plant Physiol. 110: 311-319, 1996
Soybean	FAD 2-2	Δ 12, microsomal developing seeds and vegetative tissues	18:1->18:2	no	Heppard, E.P. et al. 1996, <i>supra</i>

-5-

Borage		Δ -6	18:2 (9,12)-18:3 (6,9,12)	yes, N- terminal	Sayanova et al. Proc. Natl. Acad. Sci. USA <u>94</u> : 4211- 4216, 1997
--------	--	-------------	---------------------------------	---------------------	--

The conversion of saturated fatty acyl chains to monounsaturated species in plants appears to be confined to the chloroplasts. No Δ -9 desaturase activity has been identified in the cytoplasm or endoplasmic reticulum of plants. The soluble plant chloroplast Δ -9 desaturase is highly specific for 18:0-ACP as a substrate and does not desaturate 16:0-ACP (Somerville and Browse, *supra*). As a result, only a small amount of 16:1 is present in most higher plants, while the pool of 16:0 is concomitantly larger due to its disfavor as a substrate for the plant desaturase. By comparison, a larger amount of 18:1 is found in higher plant cells, with a correspondingly lesser amount of 18:0. Thus, for the purpose of increasing the concentration of monounsaturated lipids in a plant, the 16:0 fatty acid constitutes a significant pool of available substrate that is under-utilized by the endogenous plant desaturase.

In contrast to the plant Δ -9 desaturase, fungal and animal Δ -9 desaturases efficiently convert a wide range of saturated fatty acids with differing hydrocarbon chain lengths to monounsaturated fatty acids. The *Saccharomyces cerevisiae* enzyme, for example, efficiently desaturates even and odd chain fatty acyl CoA substrates from 13 carbons to 19 carbons in length. A broad functional homology exists among various Cyt b₅-dependent desaturases, as evidenced, for example, by the successful expression of the rat Δ -9 desaturase in yeast (Stukey et al., J. Biol. Chem. 265: 20144-20149, 1990).

-6-

The rat and yeast Δ -9 desaturase genes have been expressed in plants: both the rat and the yeast genes have been expressed in tobacco (Grayburn et al., BioTechnology 10: 675-678, 1992 (rat); Polashock et al., Plant Physiol. 100: 894-901, 1992 (yeast), and the yeast gene has also been expressed in tomato (Wang et al., J. Agric. Food Chem. 44: 3399-3402, 1996). The yeast Δ -9 desaturase has been shown to function in tobacco and tomato, leading to increases in the level of monounsaturated fatty acids (both 16:1 and 18:1) and other compounds derived from monounsaturated fatty acids (e.g., polyunsaturated fatty acids, hexanal, 1-hexanol, heptanal, trans-2-octenal) (Polashock et al., supra; Wang et al; supra). Expression of the rat desaturase also led to an increase in monounsaturated 16- and 18-carbon fatty acids (Grayburn et al., supra).

From the foregoing, it can be seen that transgenic plants expressing animal or fungal Δ -9 desaturase genes can be improved in their unsaturated fatty acid composition by virtue of the activity of the foreign enzyme. Of further advantage, it has recently been discovered that some fungal Δ -9 desaturases (e.g., *Saccharomyces cerevisiae*) are fusion proteins comprising an intrinsic Cyt b_5 domain (Mitchell & Martin, J. Biol. Chem. 270: 29766-29772, 1995). When this gene is expressed, sufficient Cyt b_5 is produced to drive the desaturase reaction at an optimum level and is not dependent on existing plant Cyt b_5 . The known animal Δ -9 desaturases do not contain this fused Cyt b_5 motif and must rely on independently-produced Cyt b_5 to provide the electrons for the reactions.

Though fungal or animal Δ -9 desaturases (e.g. the *S. cerevisiae* desaturase or the animal desaturases)

-7-

may be expressed and functional in certain plants, their expression is likely less than optimal in plants, and expression may not even be possible in other plant species, due to several factors, including differences in codon usage and codon preference in plants as compared to fungi, and among different plant species and the presence of cryptic intron splicing signals, among others. All of these factors can lead to poor expression, or no expression, of a non-plant foreign gene in a plant cell.

Accordingly, in order to make use of non-plant fatty acid desaturases, particularly those such as the *S. cerevisiae* Δ -9 desaturase comprising an internal Cyt b₅ motif, a need exists to design modified desaturase-encoding DNA molecules that are customized for expression in plant cells and specific plant tissues. It would be of even greater advantage to optimize such modified DNA molecules for expression in particular plant species, such as those that are grown and harvested primarily for oils.

SUMMARY OF THE INVENTION

According to one aspect of the invention, a synthetic fatty acid desaturase gene for expression in a multi-cellular plant is provided, the gene comprising a desaturase domain and a Cyt b₅ domain, wherein the gene is customized for expression in a plant cytoplasm. In one embodiment, the synthetic gene is customized for expression in a monocotyledonous plant. In another embodiment, the synthetic gene is customized for expression in a dicotyledonous plant. In a preferred embodiment, the synthetic gene is customized for expression in a plant genus selected from the group consisting of *Arabidopsis*, *Brassica*, *Phaseolus*, *Oryza*,

-8-

Olea, *Elaeis* (Oil Palm) and *Zea*.

In a preferred embodiment of the invention, the desaturase is a cytosolic Δ -9 desaturase. The *Saccharomyces cerevisiae* Δ -9 desaturase is particularly preferred.

In another embodiment of the invention, the synthetic gene is customized from a naturally occurring gene comprising both a desaturase domain and a cyt b_5 domain. Alternatively, the synthetic gene is a chimeric gene comprising a desaturase domain and a heterologous cyt b_5 domain.

In another embodiment, the synthetic gene is customized from a naturally occurring gene such that the synthetic gene and the naturally occurring gene encode an identical amino acid sequence. Alternatively, the synthetic gene is customized from a naturally occurring gene such that the synthetic gene and the naturally occurring gene encode a similar and functionally conserved amino acid sequence.

In another embodiment, a naturally occurring or a synthetic gene is customized so that specific amino acid modification are made to enhance the function of the encoded protein. Examples of such modifications include changing amino acids that are subjected to phosphorylation or other post-translational modifications that may alter or regulate the activity of the Δ -9 desaturase enzyme.

In another embodiment of the invention, elements of a naturally occurring or a synthetic desaturase gene that are not essential for enzymatic function are replaced or linked with elements derived from plant ER lipid biosynthetic genes that are normally expressed in maturing seeds or other plant tissues. The

-9-

improved expression of the modified gene produced by the inclusion or substitution of plant DNA sequences in the synthetic gene will result from native plant signal or control elements in those sequences that affect
5 desaturase gene expression at one or more levels.

According to another aspect of the invention, a method is provided for constructing and customizing a bifunctional desaturase/cyt b_5 encoding gene for expression in the cytosol of a multicellular plant. The
10 method comprises (a) providing a DNA molecule comprising a desaturase-encoding moiety operably linked to a cyt b_5 -encoding moiety, said DNA molecule producing the bifunctional polypeptide in a non-customized form; (b) back-translating the polypeptide sequence using preferred
15 codons for expression in a multicellular plant, thereby producing a back-translated nucleotide sequence; (c) analyzing the back-translated nucleotide sequence for features that could diminish or prevent expression in the plant cytoplasm, including, optionally (1) probable
20 intron splice sites (characterized by T-rich regions); (2) plant polyadenylation signals (e.g., AATAAA); (3) polymerase II termination sequence (e.g., CAN₍₇₋₉₎AGTNNAA, where N is any nucleotide); (4) hairpin consensus sequences (e.g., UCUUCGG); and (5) the sequence-
25 destabilizing motif ATTTA; (d) modifying the analyzed sequence to correct or remove the features that could diminish or prevent expression in the plant cytoplasm; and, optionally, (e) introducing desirable cloning features, such as restriction sites, into the sequence in
30 a manner that does not materially affect the desired codon usage or final polypeptide sequence.

The method set forth above may be adapted by incorporating into the customized gene one or more

-10-

genomic segments from plant desaturase or other ER lipid biosynthetic genes, which are determined to further optimize gene expression in plants. This method comprises (1) identifying cDNA sequences that have potential to comprise such beneficial elements, (2) creating yeast vectors expressing desaturase genes modified to contain these elements, (3) testing the vectors in a yeast expression system, (4) isolating regions from genomic DNA that are homologous to the beneficial cDNA elements, and (6) using them to construct chimeric or hybrid synthetic genes that produce functional and highly efficient desaturase activities in plant tissues.

Other features and advantages of the present invention will be better understood by reference to the drawings, detailed description and examples that follow.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1. GCG Pileup comparison of stearyl-CoA desaturase protein sequences. Sequences containing a Cyt b₅ domain are indicated with a +; sequences lacking a Cyt b₅ domain are indicated with a -; sequences still in question are indicated with a ?.

Figure 2. GCG Pileup comparison of Cytochrome b₅ protein sequences.

DETAILED DESCRIPTION OF THE INVENTION

I. Definitions

Various terms relating to the biological molecules of the present invention are used herein above and also throughout the specifications and claims.

The term "promoter region" refers to the 5'

-11-

regulatory regions of a gene.

The term "reporter gene" refers to genetic sequences which may be operably linked to a promoter region forming a transgene, such that expression of the reporter gene coding region is regulated by the promoter and expression of the transgene is readily assayed.

The term "selectable marker gene" refers to a gene product that when expressed confers a selectable phenotype, such as antibiotic resistance, on a transformed cell or plant.

The term "operably linked" means that the regulatory sequences necessary for expression of the coding sequence are placed in the DNA molecule in the appropriate positions relative to the coding sequence so as to effect expression of the coding sequence. This same definition is sometimes applied to the arrangement of coding sequences and transcription control elements (e.g. promoters, enhancers, and termination elements) in an expression vector.

The term "DNA construct" refers to genetic sequence used to transform plants and generate progeny transgenic plants. These constructs may be administered to plants in a viral or plasmid vector. Other methods of delivery such as Agrobacterium T-DNA mediated transformation and transformation using the biolistic process are also contemplated to be within the scope of the present invention. The transforming DNA may be prepared according to standard protocols such as those set forth in "Current Protocols in Molecular Biology", eds. Frederick M. Ausubel et al., John Wiley & Sons, 1999.

This invention provides synthetic DNA molecules (sometimes referred to herein as "synthetic genes") that

-12-

encode a fatty acid desaturase useful for modifying the fatty acid composition of a plant. The DNA molecules describe in accordance with this invention are superior to DNA molecules currently available for this purpose, in two important respects: (1) they encode a dual-domain polypeptide (sometimes referred to herein as a "bifunctional polypeptide or protein"), one domain being the fatty acid desaturase, and the other domain being cytochrome b_5 , a protein required to support the electron transfer events that enable the desaturase to function; and (2) they are customized for expression in the cytosol of plant cells, and further customized for expression in particular selected plant species.

Design of synthetic genes of the present invention is accomplished in two broad steps. First, the two components (the desaturase-encoding component and the Cyt b_5 -encoding component) are selected and linked together, if they do not occur together naturally. Second, the DNA molecule is optimized for expression in the cytosol of a plant cell, or further for expression in a particular plant species, or group of species.

With regard to the first step, it should be noted that several fungal, animal and plant species, including yeast, are now known to contain naturally-occurring genes encoding dual-domain cytoplasmic fatty acid desaturases. As mentioned above, the yeast and rat Δ -9 desaturase genes have been expressed and shown to function in plants. However, prior to the present invention, it was not appreciated that the bifunctional yeast desaturase offers a significant advantage over the single-function animal desaturase in plant cells, where the requisite Cyt b_5 is available only in small amounts, and the yeast protein can provide its own supply of Cyt

-13-

b₅.

With regard to the second step - optimization for expression in the plant cytosol - it was discovered in accordance with the present invention that a non-plant
5 desaturase-encoding gene, such as the yeast *OLE 1*, though expressed in some plants, may not be optimally expressed in those plants. Furthermore, the inventors have found that the yeast gene is poorly expressed in other plant species, thus highlighting the advantages obtainable by
10 optimizing such a gene for expression in a plant cell.

Sections II-IV below describe in detail how to design and use the synthetic genes of the present invention. To the extent that specific materials are mentioned, it is merely for purposes of illustration and
15 is not intended to limit the invention. Unless otherwise specified, general biochemical and molecular biological procedures, such as those set forth in Sambrook et al., Molecular Cloning, Cold Spring Harbor Laboratory (1989) (hereinafter "Sambrook et al.") or Ausubel et al. (eds)
20 Current Protocols in Molecular Biology, John Wiley & Sons (1999) (hereinafter "Ausubel et al.") are used.

II. Design and construction of the synthetic DNA molecules

A. Selection of component DNA segments

25 This invention contemplates the use of the following source DNAs, which are thereafter modified for expression in plants, if necessary:

1. naturally occurring genes or cDNAs that encode dual domain polypeptides comprising a desaturase
30 domain and a Cyt b₅ domain;
2. chimeric genes in which a desaturase-encoding sequence from one source (e.g., the desaturase domain of a dual domain fungal Δ -9 desaturase, or the

-14-

single domain rat desaturase), is linked to a Cyt b_5 -encoding sequence from a different source (e.g., a plant);

3. chimeric genes in which a sequence that
5 encodes a fragment of a naturally occurring plant Cyt b_5
(e.g. the heme binding fold, or residues that comprise
the electron donor or acceptor sites, or residues that
act as membrane targeting or retention signals, or
residues that act to stabilize the protein in the plant
10 cytoplasmic environment) is substituted for homologous
regions in the cytochrome b_5 domain of a dual domain
polypeptide such as the yeast Δ -9 desaturase; and

4. chimeric genes in which elements that encode
the essential enzymatic domains from one source (e.g. a
15 native or synthetic gene derived from a fungal Δ -9
desaturase) are linked to elements derived from native
plant desaturases that enhance transcription, mRNA
processing, mRNA stability, protein folding and
maturation, membrane targeting or retention, or protein
20 stability.

Naturally occurring genes or cDNAs that encode
dual domain desaturase/Cyt b_5 proteins have been
identified in several fungal species, including
Saccharomyces cerevisiae, *Pichia augusta*, *Histoplasma*
25 *capsulatum* and *Cryptococcus curvatus* (See Fig. 1).
Naturally occurring genes or cDNA=s that encode
independent, diffusible Cyt b_5 proteins have been
identified in several plant species, including *Nicotiana*
tabacum (tobacco), *Oryza sativa* (rice), *Cuscuta reflexa*
30 (southern Asian dodder), *Arabidopsis thaliana*, *Brassica*
oleracea and *Olea europaea* (olive). A N-terminal Cyt b_5
domain of a Δ -6 desaturase has also been identified in
the plant *Borago officinalis*, and in the *Saccharomyces*

-15-

cerevisiae FAH1 gene that encodes a very long chain fatty acid hydroxylase. Genes or cDNAs from these species, as well as DNA from any other species identified in the future as encoding such a dual domain protein, are contemplated for use in the synthetic genes of the present invention.

In a preferred embodiment, the yeast OLE1 gene is used. This embodiment is described in detail in Example 1.

The second strategy involves linking a DNA segment encoding a fatty acid desaturase from one source with a Cyt b₅ domain from another source. In a preferred embodiment, this chimeric gene is fashioned after the naturally-occurring dual function genes discussed above. That is, the Cyt b₅ domain and the desaturase domain are situated in the same positions respective to each other as is found in the naturally occurring genes (see, e.g., Mitchell & Martin, J. Biol. Chem. 270: 29766-29772, 1996).

The chimeric dual-domain proteins of the invention are prepared by recombinant DNA methods, in which DNA sequences encoding each domain are operably linked together such that upon expression, a fusion protein having the desaturase and Cyt b₅ functions described above is produced. As defined above, the term "operably linked" means that the DNA segments encoding the fusion protein are assembled with respect to each other, and with respect to an expression vector in which they are inserted, in such a manner that a functional fusion protein is effectively expressed. The selection of appropriate promoters and other 5' and 3' regulatory regions, as well as the assembly of DNA segments to form an open reading frame, employs standard methodology well

-16-

known to those skilled in the art.

Thus, preparing the chimeric DNAs of the invention involves selecting DNA sequences encoding each of the aforementioned components and operably linking the
5 respective sequences together in an appropriate vector. The sequences are thereafter expressed to produce the dual-function protein.

Genes or cDNAs that encode single-function cytoplasmic Δ -9 fatty acid desaturases have been
10 identified in a diverse array of procaryotic and eucaryotic species, including insects, fungi and mammals, but not plants (Fig. 1). Genes or cDNAs from any of these species, as well as DNA from any other species identified in the future as encoding a fatty acid
15 desaturase, are contemplated for use in the synthetic genes of the present invention.

In preferred embodiments, desaturase-encoding genes from eucaryotes, most preferably fungi or mammals, are used. In a particularly preferred embodiment, a DNA
20 encoding the rat stearoyl CoA desaturase is used. This DNA has been successfully expressed in tobacco, and accordingly is expected to be useful as part of a chimeric desaturase/Cyt b_5 gene of the present invention.

Genes or cDNAs that encode Cyt b_5 proteins have
25 also been identified in a diverse array of eucaryotic species, including insects, fungi, mammals and plants. Genes or cDNAs from any of these species, as well as DNA from any other species identified in the future as encoding a Cyt b_5 protein, are contemplated for use in the
30 synthetic genes of the present invention.

In preferred embodiments, Cyt b_5 -encoding genes or cDNAs from plants are used. These DNAs are preferred because they naturally comprise the codon usage preferred

-17-

in plants, so require little, if any, of the modification steps described below for non-plant genes. Particularly preferred, if available, are Cyt b₅-encoding DNAs from the same plant species (or group of species) to be transformed with the chimeric gene. For instance, synthetic chimeric genes constructed for transformation of *Brassica* species might comprise a stearyl CoA-encoding domain from rat and a Cyt b₅ domain from *Brassica* (see Figs. 1 and 2 for specific sources). This chimeric DNA would require optimization for expression in *Brassica* only in the desaturase domain.

With respect to the naturally-occurring dual domain-encoding genes, as well as the chimeric genes discussed above, it will be appreciated that the DNA molecules can be prepared in a variety of ways, including DNA synthesis, cloning, mutagenesis, amplification, enzymatic digestion, and similar methods, all available in the standard literature. Additionally, certain DNA molecules can be obtained by access to public repositories, such as the American Type Culture Collection. Alternatively, DNA molecules that are not readily available, and/or for which sequence information is not available, can be isolated from biological sources using standard hybridization methods and homologous probes that are available.

B. Optimization for expression in plants

The second step in designing the synthetic DNA molecules of the invention is to customize (i.e. optimize) their sequence for expression in the plant cytoplasm. This is accomplished by performing one or more of the steps listed below on the coding sequence of the above described non-plant (or chimeric)

-18-

desaturase/Cyt b₅-encoding DNA molecules.

1. From the peptide sequence encoded by the DNA, back translate using an appropriate plant codon usage table, making certain in particular that the most preferred translation termination codon is used.

2. Visually, or with the aid of computer software, analyze the back-translated nucleotide sequence for features that could diminish or prevent expression in the plant cytoplasm. Such features include: (1) probable intron splice sites (characterized by T-rich regions); (2) plant polyadenylation signals (e.g., AATAAA); (3) polymerase II termination sequence (e.g., CAN₍₇₋₉₎AGTNNA, where N is any nucleotide); (4) hairpin consensus sequences (e.g., UCUUCGG); and (5) the sequence-destabilizing motif ATTTA (Shah & Kamen, Cell 46: 659-667, 1986). These features have been described in the art (U.S. Patent No., 5,500,365 to Fischhoff et al.; U.S. Patent No. 5,380,831 to Adang et al.).

3. Modify the back-translated sequence in light of any "problem" sequences identified in step 2. Note that this step may require the introduction of codons that are not the most preferred, but instead are second or third-most preferred, in order to eliminate the more problematic sequences identified in step 2.

4. Introduce desirable cloning features, such as restriction sites, into the sequence in a manner that does not materially affect the desired codon usage or final polypeptide sequence.

The aforementioned optimization procedure can be performed so that the final polypeptide sequence is identical to the initial polypeptide sequence, even though the underlying nucleotide sequence has been modified. This is a preferred embodiment of the

-19-

invention. However, it is entirely feasible to modify the initial sequence such that the final sequence is not identical to the initial sequence, either by virtue of amino acid substitutions, insertions or deletions. The more that is known about the structure/function relationship in a particular desaturase protein, the more liberties can be taken in modifying the protein sequence during the DNA optimization process. For instance, the present inventors have shown that the entire "coiled coil" domain of the yeast *OLE1* gene can be deleted, and the protein remains functional. Thus, it appears that *OLE1* can tolerate significant modification in the encoded protein without losing its biological activity.

Codon usage tables for a variety of plants, including general plant codon usage tables, tables for dicots, tables for monocots, and tables for particular species, are widely available. Some of these are reproduced in Example 1 below. One good location to access such tables is the website:

<http://biochem.octago.ac.nz.800/Transterm/codons.html>.

In an exemplary embodiment of the present invention, the above process is applied to the coding sequence of the yeast *OLE1* gene, which encodes a cytoplasmically expressed dual-domain protein comprising a Δ -9 fatty acid desaturase domain and a Cyt b_5 domain. Optimization of the *OLE1* gene for expression in *Arabidopsis* and related species is described in detail in Example 1.

In another preferred embodiment, the coding sequence of the rat stearyl CoA desaturase is modified for expression in plants according to the methods

-20-

described above. The modified sequence is operably linked to a coding sequence for a Cyt b₅ domain, preferably from a plant, and most preferably from *Brassica*. In this regard, it has been shown that

5 expression of this rat desaturase in tobacco produces a functional protein that increases the 16:1 fatty acid content of plant tissues. Splice site prediction analysis of the rat desaturase reveals that there are no plant intron-like sequences within the open reading

10 frame. However, codon usage analysis reveals that this desaturase possesses a number of codons that are not optimal for expression in plants, particularly *Arabidopsis* or *Brassica*.

In another preferred embodiment, the protein

15 coding sequences of the modified vectors described above are further modified to increase desaturase activity. This is done by altering specific amino acids in the encoded protein that control desaturase activity through post-translational modifications. These modifications

20 are presumed to increase the level of desaturase activity in the host plant by stabilizing the desaturase protein or by increasing catalytic activity of the desaturase. Post translational modifications such as protein phosphorylation or dephosphorylation have been shown to

25 alter activity of a number of enzymes by a number of different mechanisms. These include increasing or decreasing enzyme activity or protein stability, or changing the intracellular location of the enzyme. An examination of a wide range of Δ -9 desaturase enzymes

30 reveals the existence of a number of highly conserved potential phosphorylations sites that could serve as sequences that regulate desaturase activity. These are shown in bold face on the pile-up diagram in Figure 3 and

-21-

are summarized in Table 1 of Example 3. The high degree of homology between these sites suggests that these sequences may also be recognized by host plant phosphorylating or dephosphorylating enzymes. If phosphorylation of an amino acid within one of the sites increases the activity of the desaturase, the nucleic acid sequence corresponding to that amino acid can be altered to encode a negatively charged amino acid at that site to permanently increase the activity of the protein in the host. If phosphorylation of an amino acid within the site reduces the activity of the desaturase enzyme, the nucleic acid sequence can be altered to replace that amino acid with a neutral amino acid that will permanently increase the activity of the enzyme.

In another preferred embodiment, elements of the genes in the modified vectors described above are further modified and improved by the linkage or substitution of sequences derived from native plant ER lipid biosynthetic genes. Those sequences contain elements that improve the desaturase activity by increasing the efficiency of gene expression, intracellular protein targeting and/or enzyme stability. This is done by identifying elements of the engineered desaturase gene that can be replaced or linked with elements of a plant gene without significantly affecting the desired activity or specificity of the resulting enzyme. Genes and cDNAs that encode ER lipid biosynthetic enzymes from *Brassica*, *Arabidopsis*, *Nicotiana tabacum*, Borage, maize, sunflower and soybeans, as well as similar plant genes from any other species that are identified in the future, are contemplated for use in the synthetic genes of the present invention.

In connection with the aforementioned

-22-

embodiment, but not limited thereto, it is particularly useful in many cases to pre-test constructs of the invention in a yeast expression system, in order to eliminate constructs that work poorly before taking the more labor- and time-intensive step of testing them in plants. Accordingly, this step may be incorporated into the methods described herein.

10 **III. Construction of vectors for transforming plant nuclei, and production of transgenic plants expressing synthetic genes of the invention**

 The synthetic genes of the present invention are intended for use in producing transgenic plants that optimally express a dual-function desaturase/Cyt b₅ protein in the cytoplasm of plant cells. Transformation of plant nuclei to produce transgenic plants may be accomplished according to standard methods known in the art. These include, but are not limited to, *Agrobacterium* vectors, PEG treatment of protoplasts, biolistic DNA delivery, UV laser microbeam, gemini virus vectors, calcium phosphate treatment of protoplasts, electroporation of isolated protoplasts, agitation of cell suspensions with microbeads coated with the transforming DNA, direct DNA uptake, liposome-mediated DNA uptake, and the like. Such methods have been published in the art. See, e.g., Methods for Plant Molecular Biology, Weissbach & Weissbach eds., Academic Press, Inc. (1988); Methods in Plant Molecular Biology, Schuler & Zielinski, eds., Academic Press, Inc. (1989); Plant Molecular Biology Manual, Gelvin Schilperoort, Verma, eds., Kluwer Academic Publishers, Dordrecht (1993); and Methods in Plant Molecular Biology - A Laboratory Manual, Maliga, Klessig, Cashmore, Gruissem &

-23-

Varner, eds., Cold Spring Harbor Press (1994).

The method of transformation depends upon the plant to be transformed. The biolistic DNA delivery method is useful for nuclear transformation, and is a preferred method for practice of this invention. In another embodiment of the invention, *Agrobacterium* vectors are used to advantage for efficient transformation of plant nuclei.

In a preferred embodiment, the synthetic gene is introduced into plant nuclei in *Agrobacterium* binary vectors. Such vectors include, but are not limited to, BIN19 (Bevan, Nucl. Acids Res., 12: 8711-8721, 1984) and derivatives thereof, the pBI vector series (Jefferson et al., EMBO J., 6: 3901-3907, 1987), and binary vectors pGA482 and pGA492 (An, Plant Physiol., 81: 86-91, 1986). A new series of *Agrobacterium* binary vectors, the pPZP family, is preferred for practice of the present invention. The use of this vector family for plant transformation is described by Svab et al. in Methods in Plant Molecular Biology - A Laboratory Manual, Maliga, Klessig, Cashmore, Gruissem and Varner, eds., Cold Spring Harbor Press (1994).

Using an *Agrobacterium* binary vector system for transformation, the synthetic gene of the invention is linked to a nuclear drug resistance marker, such as kanamycin or gentamycin resistance. *Agrobacterium*-mediated transformation of plant nuclei is accomplished according to the following procedure:

(1) the gene is inserted into the selected *Agrobacterium* binary vector;

(2) transformation is accomplished by co-cultivation of plant tissue (e.g., leaf discs) with a suspension of recombinant *Agrobacterium*, followed by

-24-

incubation (e.g., two days) on growth medium in the absence of the drug used as the selective medium (see, e.g., Horsch et al., Science 227: 1229-1231, 1985);

(3) plant tissue is then transferred onto the selective medium to identify transformed tissue; and

(4) identified transformants are regenerated to intact plants.

It should be recognized that the amount of expression, as well as the tissue specificity of expression of the synthetic genes in transformed plants can vary depending on the position of their insertion into the nuclear genome. Such position effects are well known in the art; see Weising et al., Ann. Rev. Genet., 22: 421-477 (1988). For this reason, several nuclear transformants should be regenerated and tested for expression of the synthetic gene.

IV. Uses of the synthetic genes and transgenic plants expressing those genes

The synthetic desaturase genes of the invention and transgenic plants expressing those genes can be used for several agriculturally beneficial purposes. For instance, they can be used in oil-producing crops (e.g., corn, soybean, sunflower, rapeseed) to increase the overall percentages of monounsaturated fatty acids in those oils, thereby improving their health-promoting qualities. In this regard, the production of transgenic rapeseed plants (*Brassica napus*) is of particular interest in this invention. Example 1 describes a synthetic yeast desaturase gene modified for expression in *Arabidopsis*. Because the codon usage of *Brassica* is very similar to that of *Arabidopsis*, it is expected that the synthetic gene described in Example 1 will be as well

-25-

expressed in *Brassica* as it is in *Arabidopsis*.

Another use for the synthetic genes of the invention is to modify the flavors of certain fruit or vegetable crops. It has already been shown that
5 expression of the un-modified yeast Δ -9 desaturase gene in tomato results in alterations in fatty acid composition and fatty acid-derived flavor compounds (Wang et al., 1996, *supra*). The synthetic, plant-optimized version of this gene is expected to function similarly,
10 and also to be more efficiently expressed in plant cells.

Another use for the synthetic genes of the invention is to facilitate the formation of omega-5 anacardic acids, a class of secondary compounds derived from the Δ -9 desaturation of 14:0 in pest-resistant
15 geraniums (Schultz et al., Proc. Natl. Acad. Sci. USA, 93: 877-885, 1996). It has been shown that formation of these compounds proceeds from the expression of Δ 9 desaturase activity resulting in the formation of Δ 9 14:1. Subsequent elongation of these molecules leads to
20 the formation of omega-5 22:1 and 24:1 in the trichome exudate that leads to pest resistance against spider mites and aphids.

Another use for the synthetic genes of the invention are in the modification of membrane lipid fatty
25 acyl composition to alter the properties of the cytoplasmic and plasma membranes of the cell. These may affect functions such membrane associated activities that are associated with membrane functions such as signal transduction, endocytosis or exocytotic events, entry of
30 fungal or viral pathogens into the cell, and temperature or environmentally caused stress that causes physical changes in the fluid properties of the plasma membrane or internal cell membranes. Plants defective in desaturases

-26-

have been reported (Somerville and Browse, *supra*). These mutant plants contain higher than normal levels of saturated fatty acids that may lower membrane fluidity under normal growing conditions. Thus the effects of temperature on these plants involved high temperature tolerance as opposed to chilling tolerance. These studies yielded interesting information that has relevance to temperature stress in general. A mutant of *Arabidopsis* deficient in 16:0 desaturation (Hugly et al, Plant Physiol. 90: 1134-1142) for example, has been shown to appear and grow normally at non-stressful temperatures. Under high temperature conditions, however, the mutant performs better than controls in growth and biosynthetic studies. Higher temperature stability was also noted in pea thylakoids following catalytic hydrogenation (Thoman et al. Biochem. Biophys. Acta 849: 131-140, 1986).

The following examples are provided to describe the invention in greater detail. They are intended to illustrate, not to limit, the invention.

EXAMPLE 1

25 Modification of the *Saccharomyces cerevisiae* OLE1 Gene for Expression in *Arabidopsis* and Related Species

When introduced into tobacco and tomato plants, the yeast Δ -9 desaturase gene (*OLE1*) was shown to desaturate palmitate and stearate, thereby reducing the levels of saturated fatty acids in triglycerides (Polashock et al., *supra*; Wang et al., *supra*). However, it was unclear whether optimum expression of the *OLE1* gene occurred in those species, and expression in other

-27-

plant species has been less than optimum. For example, the present inventors have found that the level of expression of the *OLE1* gene in tobacco (Polashock, et al., Plant Physiol. 100:894-901, 1992) and *Arabidopsis* varies in different plant tissues and is generally poor in tobacco, and *Arabidopsis* seeds. Similarly, data from other investigators indicate that expression of *OLE1* in rapeseed (*Brassica napus*) seeds is also poor (U.S. Patent No. 5,777,201, to Poutre, et al.).

10 Differential expression of heterologous genes in plants can be caused by several factors. It is often due to the presence of cryptic intron splicing signals. Thus, it is possible that the multiple banding patterns observed in northern blots of *OLE1*-transformed tobacco
15 (Polashock et al., supra) are due to splicing of the *OLE1* mRNA.

In plants, the mRNA splicing mechanism is less well defined than in mammalian or yeast systems. There is some conservation of the 5' and 3' splicing signals
20 but there is no conserved internal splice signal. However, with the accumulation of plant genomic DNA sequence data, it is now becoming possible to predict with some accuracy where intron splicing will occur (Hebsgaard, S.M., P.G. Korning, N. Tolstrup, J. Engelbrecht, P. Rouze and S. Brunak, Nucleic Acids
25 Research 24(17): 3439-3452, 1996). In fact, computer programs that predict splice sites have now been developed (the "PlantNetGene" server for splice site predictions: <http://www.cbs.dtu.dk/NetPlantGene.html>).
30 From these sources, it appears that plant introns are typically identified as T rich sequences.

Another factor affecting expression of foreign genes in plants is codon preference. It is now well

-28-

known that preference for certain codons exist among different phyla, classes, families, genera and species. Accordingly, by modifying a DNA sequence so that it uses codons preferred in a particular organism, expression of that sequence can be optimized.

Other factors affecting the expression of foreign genes in plants include the presence of putative polyadenylation signals, hairpin cleavage consensus motifs, polymerase II termination sequences and the Shaw-Kamen sequence pattern ATTTA.

This example describes the design and construction of "pl-ole1", a modified *Saccharomyces cerevisiae* OLE1 gene optimized for expression in *Arabidopsis* and other plant species.

The nucleotide sequence of the *Saccharomyces cerevisiae* OLE1 gene coding sequence has been described in U.S. Patent No. 5,057,419 to Martin et al. (incorporated by reference herein) and is set forth below for convenience as SEQ ID NO:1 (open reading frame starts at +11). The *S. cerevisiae* Δ-9 desaturase amino acid sequence encoded by OLE1 is set forth as SEQ ID NO:2.

I. Design of pl-ole1

To modify OLE1 for optimum expression in plants, the OLE1 sequence was first analyzed for cryptic plant splice signals, using the PlantNetGene server for splice site predictions. This analysis identified a number of "high confidence" intron splice signals in the OLE1 sequence. These are shown below (positions correspond to position numbers in SEQ ID NO:1).

30 Donor splice site, direct strand:

5' - 3'			5' - 3'
<u>Position</u>	<u>Strand</u>	<u>Confidence</u>	<u>exon^intron</u>
(Start ATG = +1)			

-29-

397	+	1.00	GCTCTCTCTG^GTAAAGTACC
1052	+	0.85	CTATTAAGTG^GTACCAATAC
1074	+	1.00	CCCAACTAAG^GTTATCATCT

Acceptor splice site, direct strand:

5	5' - 3'			5' - 3'
	<u>Position</u>	<u>Strand</u>	<u>Confidence</u>	<u>intron^exon</u>
	500	+	0.86	GGTCTCACAG^ATCTTACTCC

10 Next, the *OLE1* peptide sequence (SEQ ID NO:2) was back-translated using an *Arabidopsis thaliana* codon usage table, as shown below. Codon usage in *Arabidopsis* and several other plant species, including *Brassica napus*, *Phaseolus vulgaris* and *Zea mays* is very similar,

15 as can be seen by a comparison with the respective codon usage tables of those species, also shown below (the codon usage table of *Saccharomyces cerevisiae* is shown for comparison; codon usage tables taken from <http://biochem.otago.ac.nz:800/Transterm/codons.html>).

20

Arabidopsis thaliana.

	AmAcid	Codon	Number	/1000	Fraction	..
	Gly	GGG	6027.00	10.31	0.14	
25	Gly	GGA	15393.00	26.32	0.37	
	Gly	GGT	14890.00	25.46	0.35	
	Gly	GGC	5654.00	9.67	0.13	
	Glu	GAG	19825.00	33.90	0.51	
30	Glu	GAA	18672.00	31.93	0.49	
	Asp	GAT	20862.00	35.67	0.65	
	Asp	GAC	11061.00	18.91	0.35	
	Val	GTG	10414.00	17.81	0.26	
35	Val	GTA	5145.00	8.80	0.13	
	Val	GTT	16157.00	27.63	0.41	
	Val	GTC	8156.00	13.95	0.20	
	Ala	GCG	5361.00	9.17	0.13	
40	Ala	GCA	10552.00	18.04	0.25	
	Ala	GCT	18782.00	32.12	0.45	

-30-

	Ala	GCC	7249.00	12.40	0.17
	Arg	AGG	6684.00	11.43	0.22
	Arg	AGA	10280.00	17.58	0.34
5	Ser	AGT	7369.00	12.60	0.16
	Ser	AGC	6399.00	10.94	0.14
	Lys	AAG	20436.00	34.94	0.55
	Lys	AAA	16882.00	28.87	0.45
10	Asn	AAT	11658.00	19.93	0.47
	Asn	AAC	12987.00	22.21	0.53
	Met	ATG	14817.00	25.34	1.00
	Ile	ATA	6571.00	11.24	0.21
15	Ile	ATT	13028.00	22.28	0.41
	Ile	ATC	11855.00	20.27	0.38
	Thr	ACG	4346.00	7.43	0.14
	Thr	ACA	8703.00	14.88	0.28
20	Thr	ACT	10909.00	18.65	0.36
	Thr	ACC	6720.00	11.49	0.22
	Trp	TGG	6868.00	11.74	1.00
	End	TGA	652.00	1.11	0.44
25	Cys	TGT	5641.00	9.65	0.58
	Cys	TGC	4154.00	7.10	0.42
	End	TAG	252.00	0.43	0.17
	End	TAA	591.00	1.01	0.40
30	Tyr	TAT	8052.00	13.77	0.47
	Tyr	TAC	8965.00	15.33	0.53
	Leu	TTG	11727.00	20.05	0.22
	Leu	TTA	6361.00	10.88	0.12
35	Phe	TTT	11703.00	20.01	0.47
	Phe	TTC	13066.00	22.34	0.53
	Ser	TCG	4830.00	8.26	0.10
	Ser	TCA	9033.00	15.45	0.19
40	Ser	TCT	13022.00	22.27	0.28
	Ser	TCC	6214.00	10.63	0.13
	Arg	CGG	2531.00	4.33	0.08
	Arg	CGA	3142.00	5.37	0.10
45	Arg	CGT	5680.00	9.71	0.19
	Arg	CGC	2100.00	3.59	0.07
	Gln	CAG	9564.00	16.35	0.47
	Gln	CAA	10908.00	18.65	0.53

-31-

	His	CAT	7466.00	12.77	0.58
	His	CAC	5415.00	9.26	0.42
	Leu	CTG	5669.00	9.69	0.11
5	Leu	CTA	5350.00	9.15	0.10
	Leu	CTT	14395.00	24.61	0.27
	Leu	CTC	9751.00	16.67	0.18
	Pro	CCG	4676.00	8.00	0.17
10	Pro	CCA	9131.00	15.61	0.33
	Pro	CCT	10732.00	18.35	0.39
	Pro	CCC	3331.00	5.70	0.12

15 *Brassica napus*

	AmAcid	Codon	Number	/1000	Fraction	..
	Gly	GGG	730.00	11.21	0.13	
20	Gly	GGA	2042.00	31.37	0.36	
	Gly	GGT	1952.00	29.99	0.35	
	Gly	GGC	892.00	13.70	0.16	
	Glu	GAG	2119.00	32.55	0.55	
25	Glu	GAA	1764.00	27.10	0.45	
	Asp	GAT	1895.00	29.11	0.56	
	Asp	GAC	1478.00	22.70	0.44	
	Val	GTG	1231.00	18.91	0.28	
30	Val	GTA	493.00	7.57	0.11	
	Val	GTT	1624.00	24.95	0.36	
	Val	GTC	1124.00	17.27	0.25	
	Ala	GCG	615.00	9.45	0.13	
35	Ala	GCA	1167.00	17.93	0.24	
	Ala	GCT	2028.00	31.15	0.42	
	Ala	GCC	1056.00	16.22	0.22	
	Arg	AGG	697.00	10.71	0.22	
40	Arg	AGA	996.00	15.30	0.32	
	Ser	AGT	736.00	11.31	0.15	
	Ser	AGC	803.00	12.34	0.17	
	Lys	AAG	2243.00	34.46	0.55	
45	Lys	AAA	1817.00	27.91	0.45	
	Asn	AAT	1058.00	16.25	0.37	
	Asn	AAC	1811.00	27.82	0.63	
	Met	ATG	1538.00	23.63	1.00	

-32-

	Ile	ATA	669.00	10.28	0.20
	Ile	ATT	1271.00	19.52	0.37
	Ile	ATC	1461.00	22.44	0.43
5	Thr	ACG	563.00	8.65	0.15
	Thr	ACA	1059.00	16.27	0.28
	Thr	ACT	1154.00	17.73	0.30
	Thr	ACC	1073.00	16.48	0.28
10	Trp	TGG	798.00	12.26	1.00
	End	TGA	69.00	1.06	0.37
	Cys	TGT	517.00	7.94	0.50
	Cys	TGC	509.00	7.82	0.50
15	End	TAG	33.00	0.51	0.18
	End	TAA	83.00	1.28	0.45
	Tyr	TAT	792.00	12.17	0.38
	Tyr	TAC	1283.00	19.71	0.62
20	Leu	TTG	1051.00	16.14	0.20
	Leu	TTA	508.00	7.80	0.09
	Phe	TTT	1003.00	15.41	0.39
	Phe	TTC	1562.00	23.99	0.61
25	Ser	TCG	475.00	7.30	0.10
	Ser	TCA	856.00	13.15	0.18
	Ser	TCT	1147.00	17.62	0.24
	Ser	TCC	799.00	12.27	0.17
	Arg	CGG	219.00	3.36	0.07
30	Arg	CGA	297.00	4.56	0.09
	Arg	CGT	659.00	10.12	0.21
	Arg	CGC	275.00	4.22	0.09
35	Gln	CAG	1188.00	18.25	0.50
	Gln	CAA	1168.00	17.94	0.50
	His	CAT	651.00	10.00	0.49
	His	CAC	672.00	10.32	0.51
40	Leu	CTG	592.00	9.09	0.11
	Leu	CTA	579.00	8.89	0.11
	Leu	CTT	1416.00	21.75	0.26
	Leu	CTC	1208.00	18.56	0.23
45	Pro	CCG	542.00	8.33	0.15
	Pro	CCA	1180.00	18.13	0.33
	Pro	CCT	1281.00	19.68	0.36
	Pro	CCC	527.00	8.10	0.15

-33-

Phaseolus vulgaris

5	Gly	GGG	371.00	13.30	0.15
	Gly	GGA	771.00	27.64	0.32
	Gly	GGT	817.00	29.29	0.34
	Gly	GGC	441.00	15.81	0.18
10	Glu	GAG	912.00	32.69	0.54
	Glu	GAA	767.00	27.50	0.46
	Asp	GAT	776.00	27.82	0.55
	Asp	GAC	625.00	22.41	0.45
15	Val	GTG	661.00	23.70	0.36
	Val	GTA	174.00	6.24	0.09
	Val	GTT	653.00	23.41	0.36
	Val	GTC	346.00	12.40	0.19
20	Ala	GCG	180.00	6.45	0.09
	Ala	GCA	528.00	18.93	0.26
	Ala	GCT	791.00	28.36	0.39
	Ala	GCC	553.00	19.82	0.27
25	Arg	AGG	324.00	11.61	0.29
	Arg	AGA	325.00	11.65	0.29
	Ser	AGT	317.00	11.36	0.14
	Ser	AGC	353.00	12.65	0.15
30	Lys	AAG	1054.00	37.78	0.60
	Lys	AAA	697.00	24.99	0.40
	Asn	AAT	555.00	19.90	0.42
	Asn	AAC	782.00	28.03	0.58
35	Met	ATG	567.00	20.33	1.00
	Ile	ATA	274.00	9.82	0.20
	Ile	ATT	539.00	19.32	0.40
	Ile	ATC	548.00	19.65	0.40
40	Thr	ACG	166.00	5.95	0.11
	Thr	ACA	362.00	12.98	0.24
	Thr	ACT	480.00	17.21	0.32
	Thr	ACC	490.00	17.57	0.33
45	Trp	TGG	342.00	12.26	1.00
	End	TGA	34.00	1.22	0.44
	Cys	TGT	145.00	5.20	0.39
	Cys	TGC	229.00	8.21	0.61
	End	TAG	22.00	0.79	0.28
	End	TAA	22.00	0.79	0.28

-34-

	Tyr	TAT	400.00	14.34	0.40
	Tyr	TAC	597.00	21.40	0.60
	Leu	TTG	543.00	19.47	0.24
5	Leu	TTA	184.00	6.60	0.08
	Phe	TTT	458.00	16.42	0.43
	Phe	TTC	601.00	21.55	0.57
	Ser	TCG	149.00	5.34	0.06
10	Ser	TCA	416.00	14.91	0.18
	Ser	TCT	606.00	21.72	0.26
	Ser	TCC	501.00	17.96	0.21
	Arg	CGG	71.00	2.55	0.06
15	Arg	CGA	76.00	2.72	0.07
	Arg	CGT	169.00	6.06	0.15
	Arg	CGC	158.00	5.66	0.14
	Gln	CAG	437.00	15.67	0.48
20	Gln	CAA	470.00	16.85	0.52
	His	CAT	298.00	10.68	0.46
	His	CAC	355.00	12.73	0.54
	Leu	CTG	351.00	12.58	0.15
25	Leu	CTA	184.00	6.60	0.08
	Leu	CTT	569.00	20.40	0.25
	Leu	CTC	452.00	16.20	0.20
	Pro	CCG	147.00	5.27	0.08
30	Pro	CCA	694.00	24.88	0.37
	Pro	CCT	664.00	23.80	0.36
	Pro	CCC	352.00	12.62	0.19

35 *Zea mays*

	AmAcid	Codon	Number	/1000	Fraction	..
	Gly	GGG	2466.00	15.07	0.19	
40	Gly	GGA	2186.00	13.36	0.17	
	Gly	GGT	2607.00	15.93	0.20	
	Gly	GGC	5499.00	33.61	0.43	
	Glu	GAG	7364.00	45.01	0.72	
45	Glu	GAA	2823.00	17.25	0.28	
	Asp	GAT	3425.00	20.93	0.37	
	Asp	GAC	5740.00	35.08	0.63	
	Val	GTG	4365.00	26.68	0.38	

-35-

	Val	GTA	916.00	5.60	0.08
	Val	GTT	2516.00	15.38	0.22
	Val	GTC	3644.00	22.27	0.32
5	Ala	GCG	3698.00	22.60	0.24
	Ala	GCA	2517.00	15.38	0.16
	Ala	GCT	3602.00	22.01	0.24
	Ala	GCC	5481.00	33.50	0.36
10	Arg	AGG	2500.00	15.28	0.27
	Arg	AGA	1199.00	7.33	0.13
	Ser	AGT	1170.00	7.15	0.10
	Ser	AGC	2776.00	16.97	0.24
15	Lys	AAG	7241.00	44.25	0.79
	Lys	AAA	1969.00	12.03	0.21
	Asn	AAT	1946.00	11.89	0.33
	Asn	AAC	3939.00	24.07	0.67
20	Met	ATG	4071.00	24.88	1.00
	Ile	ATA	1014.00	6.20	0.13
	Ile	ATT	2099.00	12.83	0.28
	Ile	ATC	4403.00	26.91	0.59
25	Thr	ACG	1890.00	11.55	0.22
	Thr	ACA	1620.00	9.90	0.19
	Thr	ACT	1757.00	10.74	0.21
	Thr	ACC	3236.00	19.78	0.38
30	Trp	TGG	1994.00	12.19	1.00
	End	TGA	199.00	1.22	0.45
	Cys	TGT	770.00	4.71	0.28
	Cys	TGC	1963.00	12.00	0.72
35	End	TAG	121.00	0.74	0.28
	End	TAA	120.00	0.73	0.27
	Tyr	TAT	1303.00	7.96	0.27
	Tyr	TAC	3440.00	21.02	0.73
40	Leu	TTG	1807.00	11.04	0.13
	Leu	TTA	582.00	3.56	0.04
	Phe	TTT	1697.00	10.37	0.29
	Phe	TTC	4082.00	24.95	0.71
45	Ser	TCG	1620.00	9.90	0.14
	Ser	TCA	1592.00	9.73	0.14
	Ser	TCT	1792.00	10.95	0.15
	Ser	TCC	2746.00	16.78	0.23

-36-

	Arg	CGG	1505.00	9.20	0.16
	Arg	CGA	610.00	3.73	0.06
	Arg	CGT	1018.00	6.22	0.11
	Arg	CGC	2562.00	15.66	0.27
5	Gln	CAG	4280.00	26.16	0.72
	Gln	CAA	1626.00	9.94	0.28
	His	CAT	1378.00	8.42	0.36
	His	CAC	2431.00	14.86	0.64
10	Leu	CTG	4069.00	24.87	0.29
	Leu	CTA	904.00	5.52	0.07
	Leu	CTT	2415.00	14.76	0.17
	Leu	CTC	4079.00	24.93	0.29
15	Pro	CCG	2642.00	16.15	0.29
	Pro	CCA	2152.00	13.15	0.23
	Pro	CCT	2102.00	12.85	0.23
	Pro	CCC	2344.00	14.33	0.25
20					

Saccharomyces cerevisiae

	AmAcid	Codon	Number	/1000	Fraction	..
25	Gly	GGG	18129.00	6.18	0.12	
	Gly	GGA	32850.00	11.20	0.22	
	Gly	GGT	66575.00	22.69	0.45	
	Gly	GGC	28821.00	9.82	0.20	
30	Glu	GAG	57100.00	19.46	0.30	
	Glu	GAA	133513.00	45.51	0.70	
	Asp	GAT	111120.00	37.88	0.65	
	Asp	GAC	58642.00	19.99	0.35	
35	Val	GTG	32144.00	10.96	0.20	
	Val	GTA	35470.00	12.09	0.22	
	Val	GTT	63678.00	21.71	0.39	
	Val	GTC	33136.00	11.30	0.20	
40	Ala	GCG	18402.00	6.27	0.11	
	Ala	GCA	47728.00	16.27	0.30	
	Ala	GCT	58916.00	20.08	0.37	
	Ala	GCC	35917.00	12.24	0.22	
45	Arg	AGG	27990.00	9.54	0.21	
	Arg	AGA	61524.00	20.97	0.47	
	Ser	AGT	42499.00	14.49	0.16	
	Ser	AGC	29298.00	9.99	0.11	

-37-

	Lys	AAG	89539.00	30.52	0.42
	Lys	AAA	124327.00	42.38	0.58
	Asn	AAT	106379.00	36.26	0.60
	Asn	AAC	71659.00	24.43	0.40
5					
	Met	ATG	61216.00	20.87	1.00
	Ile	ATA	53773.00	18.33	0.28
	Ile	ATT	88869.00	30.29	0.46
	Ile	ATC	49422.00	16.85	0.26
10					
	Thr	ACG	24131.00	8.23	0.14
	Thr	ACA	52363.00	17.85	0.31
	Thr	ACT	58260.00	19.86	0.34
	Thr	ACC	35998.00	12.27	0.21
15					
	Trp	TGG	30707.00	10.47	1.00
	End	TGA	1901.00	0.65	0.30
	Cys	TGT	23942.00	8.16	0.62
	Cys	TGC	14448.00	4.93	0.38
20					
	End	TAG	1421.00	0.48	0.23
	End	TAA	2985.00	1.02	0.47
	Tyr	TAT	55441.00	18.90	0.57
	Tyr	TAC	42016.00	14.32	0.43
25					
	Leu	TTG	79248.00	27.01	0.28
	Leu	TTA	77691.00	26.48	0.28
	Phe	TTT	78451.00	26.74	0.59
	Phe	TTC	53809.00	18.34	0.41
30					
	Ser	TCG	25856.00	8.81	0.10
	Ser	TCA	55962.00	19.08	0.21
	Ser	TCT	69019.00	23.53	0.26
	Ser	TCC	41460.00	14.13	0.16
35					
	Arg	CGG	5414.00	1.85	0.04
	Arg	CGA	9166.00	3.12	0.07
	Arg	CGT	18429.00	6.28	0.14
	Arg	CGC	7924.00	2.70	0.06
40					
	Gln	CAG	36018.00	12.28	0.31
	Gln	CAA	78385.00	26.72	0.69
	His	CAT	40211.00	13.71	0.64
	His	CAC	22609.00	7.71	0.36
45					
	Leu	CTG	31503.00	10.74	0.11
	Leu	CTA	39789.00	13.56	0.14
	Leu	CTT	36697.00	12.51	0.13
	Leu	CTC	16401.00	5.59	0.06

-38-

Pro	CCG	15796.00	5.38	0.12
Pro	CCA	51725.00	17.63	0.41
Pro	CCT	39402.00	13.43	0.31
Pro	CCC	20387.00	6.95	0.16

5

For each amino acid, the new *pl-ole1* gene was designed the codon most preferred in *Arabidopsis*, with the following exceptions:

- 10 1. The codon for glutamine CAG was switched to CAA. Though the codon preference for glutamine is the same for both CAG and CAA in *Arabidopsis*, CAA was used since the AG motif is part of the 3' intron splice signal.
- 15 2. In *OLE1*, there are regions of high leucine/valine amino acid usage (e.g., between positions 322 to 571 of the nucleotide sequence are codons coding for 11 leucines and 7 valines). These regions correspond to the *OLE1* protein transmembrane domains. If the most
20 preferred codons in *Arabidopsis* (CTT and GTT, respectively) were used, the region would take on the characteristics of a plant intron, i.e., high T content, thereby introducing a number of highly probable 5' splice sites, which could not be removed without altering the
25 amino acid sequence. Accordingly, a mixture of alternative codons was used for these amino acids. Similar changes were also applied to two other regions of *OLE1* (positions 781 to 900 and positions 1081 to 1140).

Next, a search for problematic sequences, such
30 as putative polyadenylation signals, hairpin cleavage consensus motifs, ATTTA motifs or concatamers thereof, was conducted. Such sequences are described in detail in U.S. Patent No. 5,380,831 to Adang et al. (incorporated by reference herein). This search identified one hairpin
35 cleavage consensus motif, CTTCGG, at position 553-559 of

-39-

SEQ ID NO:1, which was removed by changing TTC to TTT (both encoding phenylalanine).

Next, a BamHI site and translation initiation consensus were added to the 5' end of the *OLE1* coding sequence (M. Kozak, J. Biol. Chem. 266(30): 19867-19870, 1991). An XbaI and a BamHI site were added to the 3' end of the coding sequence. A PacI site was introduced into the same position as the original *S. cerevisiae* *OLE1* PacI site (within the cytochrome *b₅* domain), in order to provide a convenient restriction site for construction of this and other synthetic *OLE1* genes. Other convenient restriction sites, which enable modular construction of synthetic *OLE1* genes, are inherent within the final sequence of the new *pl-ole1* gene.

Finally, the termination codon was checked against a stop codon consensus database, "TransTerm" (Dalphin et al., Nucl. Acids Res. 25(1): 246-247, 1997). The existing termination sequence, TGAT, appeared suitable for use in *Arabidopsis*, and so was not altered.

II. Construction of *pl-ole1*:

The rebuilt *pl-ole1* nucleotide sequence was constructed commercially (Operon Technologies, Inc.). The plasmid containing the rebuilt gene was designated pAMCM013. The *pl-ole1* nucleotide sequence is set forth below as SEQ ID NO:3 (open reading frame starts at +11). This sequence encodes SEQ ID NO:2, but differs from the *S. cerevisiae* *OLE1* gene (SEQ ID NO:1) in the following respects (summarized from above):

1. *Arabidopsis thaliana* codon usage; CAG switched to CAA for glutamine;
2. Translation initiation consensus added;
3. Hairpin removed;
4. Several (but not all) PlantNetGene

-40-

predicted splice sites removed;

5. Eleven leucines changed from CTT to CTC,
and 7 valines changed from GTT to GTG in positions 322-
571, which corresponds to a plant intron-like region;
5 similar changes made in regions 781-900 and 1081-1140;
valine at position 432 retained as GTT to maintain
Psp1406I site;

6. Certain leucine and valine codons were
altered so that the same codons would not appear adjacent
10 to others;

7. Intron acceptor site at position 1047
altered;

8. Restriction sites added to allow modular
construction; Psp1406I site removed at position 1441; and

15 9. PacI site introduced at position 1362; an
introduced NgoMI site at position 867 removed.

A gap alignment of SEQ ID NO: 1 (top) and SEQ
ID NO: 3 (bottom) is shown below:

Gap alignment of wild type and rebuilt OLE1 sequences.
Percent Similarity: 79.871 Percent Identity: 79.871

```

      1 TACAACAAAGATGCCAACTTCTGGAAGTACTATTGAATTGATTGACGACC 50
          ||| |||| |||| |||| |||| |||| |||| |||| |||| ||||
      1 ggatccaacaATGCCTACTTCTGGAAGTACTATCGAGCTTATCGATGATC 50

      51 AATTTCCAAAGGATGACTCTGCCAGCAGTGGCATTGTCGACGAAGTCGAC 100
          |||| || |||| |||| |||| |||| |||| |||| |||| ||||
      51 AATTCCTAAGGATGATTCTGCTTCTTCTGGAATCGTTGATGAGGTTGAT 100

     101 TTAACGGAAGCTAATATTTTGGCTACTGGTTTGAATAAGAAAGCACCAAG 150
          || || || |||| || || |||| |||| || || |||| || || ||
     101 CTTACTGAGGCTAACATCCTTGCTACTGGACTTAACAAGAAGGCTCCTAG 150

     151 AATTGTCAACGGTTTTGGTTCTTTAATGGGCTCCAAGGAAATGGTTTCCG 200
          ||| || |||| || || || || || || || || |||| |||| ||
     151 AATCGTTAACGGATTCCGATCTCTTATGGGATCTAAGGAGATGGTTTCTG 200

     201 TGGAATTCGACAAGAAGGGAAACGAAAAGAAGTCCAATTTGGATCGTCTG 250
          || || |||| |||| |||| |||| |||| || || || || || ||
     201 TTGAGTTCGATAAGAAGGGAAACGAGAAGAAGTCTAACCTTGATAGACTT 250

```


251	CTAGAAAAGGACAACCAAGAAAAAGAAAGACTAAAATTCACAT	300
251	CTTGAGAAGGATAACCAAGAGAAGGAGGCTAAGACTAAGATCCATAT	300
301	CTCCGAACAACCATGGACTTTGAATAACTGGCACCAACATTTGAACTGGT	350
301	CTCTGAGCAACCTTGGACTCTcAACAACTGGCATCAACATCTcAACTGGC	350
351	TGAACATGGTTCTTGTGGTATGCCAATGATTGGTTGGTACTTCGCT	400
351	TcAACATGGTgCTcGTcTGTGGAATGCCTATGATCGGATGGTACTTCGCT	400
401	CTCTCTGGTAAAGTACCTTTGCATTTAAACGTTTTCTCTCCGTTTT	450
401	CTcTCTGGAAAaGTgCCTCTcCATCTcAACGTTTTCcTcTTCTCTGTcTT	450
451	CTACTACGCTGTCGGTGGTGTCTATTACTGCCGTTACCATAGATTAT	500
451	CTACTACGCTGTTGGAGGAGTgTCTATCACTGCTGGATACCATAGACTcT	500
501	GGTCTCACAGATCTTACTCCGCTCACTGGCCATTGAGATTATTCTACGCT	550
501	GGTCTCATAGATCTTACTCTGCTCATTGGCCTCTTAGACTcTTCTACGCT	550
551	ATCTTCGGTTGTGCTTCCGTTGAAGGGTCCGCTAAATGGTGGGGCCACTC	600
551	ATCTTtGGATGTGCTTCTGTTGAGGGATCTGCTAAGTGGTGGGGACATTC	600
601	TCACAGAATTCACCATCGTTACACTGATACCTTGAGAGATCCTTATGACG	650
601	TCATAGAATCCATCATAGATACACTGATACTCTTAGAGATCCTTACGATG	650
651	CTCGTAGAGGTCTATGGTACTCCACATGGGATGGATGCTTTTGAAGCCA	700
651	CTAGAAGAGGACTTTGGTACTCTCATATGGGATGGATGCTTCTTAAGCCT	700
701	AATCCAAAATACAAGGCTAGAGCTGATATTACCGATATGACTGATGATTG	750
701	AACCCTAAGTACAAGGCTAGAGCTGATATCACTGATATGACTGATGATTG	750
751	GACCATTAGATTCCAACACAGACACTACATCTTGTTGATGTTATTAACCG	800
751	GACTATCAGATTCCAACATAGACATTACATCTgCTcATGCTcCTTACTG	800
801	CTTTCGTCATTCCAACCTTATCTGTGGTTACTTTTTCAACGACTATATG	850
801	CTTTCGTCaATCCCTACTCTcATCTGTGGATACTTCTTCAACGATTACATG	850


```

1451 ATGAGAGTTGCTGTTATCAAGGAGTCTAAGAAGTCTGCTATCAGAATGGC 1500
1501 TAGTAAGAGAGGTTGAAATCTACGAAAGTGGTAAGTTCTTTTAAGCATCAC 1550
1501 TTCTAAGAGAGGAGAGATCTACGAGACTGGAAAGTTCTTCTGAtctagag 1550
1551 ATTAC 1555
1551 qatcc 1555

```

An alternative version of *pl-ole1*, referred to herein as *pl-ole1-2*, was also constructed. This synthetic gene was modified only in specific codons identified as high frequency splicing signals. It was discovered that this construct is expressed equally as well as *pl-ole1* in *Arabidopsis*.

A modification of a transformation protocol of Pam Green (<http://www.bch.msu.edu/pamgreen/vac.html>) was used for the transformation of *A. thaliana* with *pl-ole1*.

-44-

The protocol was adapted from protocols by Nicole Bechtold and Andrew Bent. This protocol gives very good results, with 95% of all infiltrated plants giving rise to transformants, and a transformant in up to 1 in 25 seeds.

PROTOCOL:

1. Seeds of *Arabidopsis thaliana* ecotype Columbia were sown in lightweight plastic pots prepared in the following way: mound *Arabidopsis* soil mixture into 3 to 4 inch pots, saturate soil with *Arabidopsis* fertilizer, add more soil so that it is rounded about 0.5 above the edge.

2. Plants were grown under conditions of 16 hours light / 8 hours dark at 20°C, fertilizing with *Arabidopsis* fertilizer once a week from below, adding about 0.5 L to each flat. After 4-6 weeks, plants were considered ready for vacuum infiltration when primary inflorescence was 10-15 cm tall and the secondary inflorescences appeared at the rosette. The bolts were clipped back and 2 to 3 days was allowed for them to regrow before infiltration.

3. In the meantime, the construct was transformed into *Agrobacterium tumefaciens* strain (LBA4404). When plants were ready to transform, a 50 mL culture of LB medium containing 50 mg/L kanamycin and 50 mg/L of streptomycin was inoculated with a 1 mL overnight starter culture.

4. Cultures were grown overnight at 28° C with shaking. The culture was pelleted, the supernatant removed, and the pellet resuspended in 250 ml of infiltration medium to OD600 >0.8. Infiltration medium (1 liter) comprised 2.2 g MS salts, 1 X B5 vitamins, 50 g

-45-

sucrose, 0.5 g MES, pH to 5.7 with KOH, 0.044 % B5M
benzylaminopurine, 200 mg BSL Silwet L-77 (OSI
Specialties).

5. The resuspended culture was placed in a
5 magenta jar inside a large bell jar. Pots containing
plants to be infiltrated were inverted into the solution
so that the entire plant was covered, including rosette,
but none of the soil was submerged.

6. A vacuum of 400 mm Hg (about 17 inches) was
10 drawn. Once the vacuum level was reached, the suction was
closed and the plants allowed to remain under vacuum for
five minutes. The vacuum was then quickly released. The
pots were briefly drained, then placed on
their sides in a tray, which was covered with a humidome
15 to maintain humidity. The next day, the plants were
removed to the growth room, the pots uncovered and set
upright. Plants infiltrated with different constructs
were kept separated in different trays thereafter.

7. Plants were allowed to grow under the same
20 conditions as before. Plants were staked individually as
the bolts grew. When plants were finished flowering,
water was gradually reduced, then eliminated to allow the
plants to dry out. Seeds were harvested from each plant
individually.

8. Large selection plates were prepared: 4.3
25 g/L MS salts; 1 X B5 vitamins (optional); 1 % sucrose;
0.5 g/L MES pH to 5.7 with KOH; 0.8% phytagar -
Autoclaved, then added antibiotics (35 µg/mL kanamycin
and 250 µg/mL of carbenicillin) and 150 X 15 mm plates
30 were poured.

9. Plates were dried well in the sterile hood
before plating - 20-30 minutes with the lids open was
usually sufficient.

-46-

10. For each plant, up to 100 μ L of seeds (approximately 2500 seeds) was sterilized and plated out individually. Seeds were sterilized as follows: 1 min in 70% ethanol, 7 minutes in 50% bleach / 0.02 % Triton X-100 with vortexing, 6 rinses in sterile distilled water. Seeds were resuspended in 2 mL sterile 0.1% agarose and poured onto large selection plates as if plating phage. Plates were tilted so seeds were evenly distributed, and allowed to sit 10-15 minutes, during which time the liquid soaked into the medium. Plates were sealed with Parafilm and placed in a growth room.

11. After 7 to 10 days, transformants were visible as dark green plants. These were transferred onto "hard selection" plates (100 x 15 mm plates with same recipe as selection plates but with 1.5 % phytagar) to eliminate any pseudo-resistants, then replaced in the growth room.

12. After 10 to 14 days, the plants possessed at least two sets of true leaves. At this point, plants were transferred to soil, covered with plastic, and moved to a growth chamber with normal conditions. They were typically kept covered for several days.

References:

- 25 Bechtold N, Ellis J, Pelletier G (1998) Methods Mol Biol. 82: 259-266.
- Bent A, Kunkel BN, Dahlbeck D, Brown KL, Schmidt R, Giraudat J, Leung J, Staskawicz BJ (1994) Science 265: 1856-1860.
- 30 Koncz C, Schell J (1986) Mol. Gen. Genet. 204: 383-396.

Solutions:

-47-

1000X B5 vitamins (10 mL):

1000 mg myo-inositol

100 mg thiamine-HCl

10 mg nicotinic acid

5 10 mg pyridoxine-HCl

Dissolve in ddH₂O and store at -20 °C.

Arabidopsis fertilizer (10 liters):

50 mL 1M KNO₃

10 25 mL 1M KPO₄ (pH 5.5)

20 mL 1M MgSO₄

20 mL 1M Ca(NO₃)₂

5 mL 0.1M Fe.EDTA

10 mL micronutrients (see below)

15 Dissolve in ddH₂O and store at room temperature

Arabidopsis micronutrients (500 mL):

70 mL 0.5M boric acid

14 mL 0.5M MnCl₂

20 2.5 mL 1M CuSO₄

1 mL 0.5M ZnSO₄

1 mL 0.1M NaMoO₄

1 mL 5M NaCl

0.05 mL 0.1M CuCl₂

25 Dissolve in ddH₂O and store at room temperature

EXAMPLE 3

Customizing *OLE1* to Express

Post-Translational Modifications

30

After determining the optimized codon preferences of *OLE1* mRNA (or mRNA derived from another fungal or animal desaturase) for high level expression in the host plant, specific amino acids that are involved in

-48-

the post-translational control of enzyme activity or stability are altered to maximize the catalytic activity of the expressed enzyme. There are a number of protein kinase and/or phosphorylase consensus sequences that are highly conserved in the fungal and animal desaturases. These are shown below. First is shown a table of aligned potential phosphorylation sites in desaturases. Next is shown a pileup of Δ -9 fatty acid desaturases. PROSITE analysis of these desaturases predicts a number of potential phosphorylation sites, highlighted by bold underlined characters.

Table 2 - Aligned Potential phosphorylation sites in desaturases. Phosphorylation sites are indicated with respect to amino acid positions in the Olefp protein coding sequence. Abbreviations, S = serine, T= Threonine, A=Alanine, PKC - protein kinase C like; CK-2 - casein kinase II like; CAMP - CAMP activated kinase (PKA) like; np., not predicted as a phosphorylation site.

Residue position	Olefp Residue	Olefp sequence	Phosphorylation type	Fungal/b5	Animal	Insect
166	S	SHR	PKC	mixed, S or A	all S, SHR	N.P., A
169	T	np	PKC	S or A	all TYK or SYK	SYK
191	S	SAK	PKC	all S	all D	all A
206	T	np	CK-2	all T	all S	all S
208	T	TLRD	CK-2	all T	all T	all T
215	A	np	PKC	A or V	all S	all A
323	T	TPRD	CK-2	T or S	S or np	np
351	R	np	CK-2	all R	all S	all K
383	S	KKFS	CAMP	S or np	all S	S but np

-50-

Fileup of Δ -9 fatty acid desaturases showing potential phosphorylation sites:

	1				50
Rat	-----	-----	-----	-----MPAHM	LQE.ISSSY.
Mouse	-----	-----	-----	-----MPAHM	LQE.ISSSY.
Sheep	-----	-----	-----	-----MPAHL	LQEEISSSY.
Pig	-----	-----	-----	-----	-----SSY.
Human	-----	-----	-----	-----MPAHL	LQDDISSSY.
Hamster	-----	-----	-----	-----MPGHL	LQEEMTSSYT
Drosophila	-----	-----	-----	-----MPP	NAQAGAQSIS
Moth	-----	-----	-----	-----	-----
C.elegans	-----	-----	-----	-----MTVKTRSN	IAKKIEKDGG
S.cerevisiae	MPTSGTTIEL	IDDQFPKDDS	ASSGIVDEVD	LTEANILATG	LNKKAPRIVN
P. angusta	-----	-----	-----	-----	-----
H. capsulatum	-----	-----	-----	-----	-----
M. rouxii	-----	-----	-----	-----	-----
C. curvatus	-----	-----	-----	-----	-----
C. merolae	-----	-----	-----	MTAKVESKVR	EEEKGSNPST
	51				100
Rat	TTTTTITEPP	SGNLQNGREK	MKKVPLYLEE	DI.....RPE	MREDIHDPSY
Mouse	TTTTTITAPP	SG...NEREK	VKTVPPLHLE	DI.....RPE	MKEDIHDPTY
Sheep	TTTTTITAPP	SRVLQNGGGK	LEKTPLYLEE	DI.....RPE	MRDDIYDPNY
Pig	TTTTTITAPS	SRVLQNGGGK	SEKTPQYVEE	DI.....RPE	MKDDIYDPTY
Human	TTTTTITAPP	PGVLQNGGDK	LETMPLYLED	DI.....RPD	IKDDIYDPTY
Hamster	TTTTTITEPP	SESLO.....	.KTVPLYLEE	DI.....RPE	MKEDIYDPSY
Drosophila	DSLIAAASAA	ADAGQSPTKL	QEDSTGVLFE	CD.....VET	TDGGLVKDIT
Moth	-----	-----MPPQG	QTGGSWVLYE	TD.....AVN	TDTD..APVI
C.elegans	PETQYLAVDP	NEIIQLQEE	KKVVPKCLPA	RLPTAACKAS	QENGECQKIV
S.cerevisiae	GFGSLMGSK	MVSVEFDKKG	NEKKSNDRL	LEKDNQEKKE	AKTKIH.ISE
P. angusta	-----MGTKS	MTDVTAEEL.	..SKDSVAMM	LAKDRELKNK	YLKQKH.ISE
H. capsulatum	-----MA	LNEAPTASPV	AETAAGGKDV	VTDAARRPNS	EPKKVH.ITD
M. rouxii	-----	-----MSN	IATLTSTART	KTESMKPPLP	KTKMPP.LFD
C. curvatus	~MSASTKQAS	TTVAQPSGKP	VTNVIDPERD	DFIVPDNYVT	RTVENM.KML
C. merolae	AAADDSGAVI	PTLKPRPKPA	VEPLEREGVE	FDPQRGLVFE	KTRSSKWMSE
	101				150
Rat	QDEEGPPPKL	EYVWRNIILM	ALLHVGALYG	ITL.IPSSKV	YTLLWGIFYF
Mouse	QDEEGPPPKL	EYVWRNIILM	VLLHLGGLYG	IIL.VPSCKL	YTALFGIFYF
Sheep	QDKEGPKPKL	EYVWRNIILM	GLLHLGALYG	ITL.IPTCKI	YTFLWVLFYF
Pig	QDKEGPQKQL	EYVWRNIILM	SLLHLGALYG	IIL.IPTCKI	YTLLWAFAYF
Human	KDKEGPSPKV	EYVWRNIILM	SLLHLGALYG	ITL.IPTCKF	YTWLWGVFYY
Hamster	QDEEGPPPKL	EYVWRNIILM	ALLHLGALYG	LVL.VPSSKV	YTLLWAFVYY
Drosophila	VMKKAERLL	KLVRNIIAF	GYLHLAALYG	AYLMVTSKAW	QTCILAYFLY
Moth	VPPSAEKREW	KIVWRNVILM	GMLHIGGVYG	AYLFLTKAMW	LTDLFAFFLY
C.elegans	FLEIVIPYKM	EIVWRNVALF	AALHFAAAIG	LYQLIFEAKW	QTVIFTFLLY
S.cerevisiae	QPWTLNNWHQ	HLNWLNMVLV	CGMPMIGWYF	ALSGKVPLHL	NVFLFSVFYY
P. angusta	QPWTWENWHR	HINWLNFIIV	LAVPFAG..L	ISTKWVPLKL	HTFVTAVILY
H. capsulatum	TPITLANWHK	HISWLNVTIL	IAIPIYG..L	VOAYWVPLHL	KTALWAVVYY
M. rouxii	QPVTSKNWT	FVNWPQAILL	CVTPLIALYG	IFT..TELTK	KTLIWSWIYY
C. curvatus	PPVTRNLHK	NIQWISFLAL	TIPPAMAIYG	LCT..VPVQT	KTFIWSVYY
C. merolae	KELNELPLLQ	RINWLS.TSI	IFTPLIGT.L	IGIWFVPLQR	KTLVLAIVTY

-51-

	151				200
Rat	LISALGITAG	AHRLW <u>SHRTY</u>	KARLPLRIFL	IIANTMAFQN	DVYEWARDHR
Mouse	MTSALGITAG	AHRLW <u>SHRTY</u>	KARLPLRIFL	IIANTMAFQN	DVYEWARDHR
Sheep	VISALGITAG	VHRLW <u>SHRTY</u>	KARLPLRVFL	IIANTMAFQN	DVFEWSRDHR
Pig	LLSAVGVTAG	AHRLW <u>SHRTY</u>	KARLPLRVFL	IIANTMAFQN	DVYEWARDHR
Human	FVSALGITAG	AHRLW <u>SHRSY</u>	KARLPLRLFL	IIANTMAFQN	DVYEWARDHR
Hamster	VISIEGIGAG	VHRLW <u>SHRTY</u>	KARLPLRIFL	IIANTMAFQN	DVYEWARDHR
Drosophila	VISGLGITAG	AHRLWA <u>HRSY</u>	KAKWPLRVIL	VIFNTIAFQD	AAYHWARDHR
Moth	LCSGLGITAG	AHRLWA <u>HKS</u>	KARLPLRLLL	TLFNTIAFQD	AVIDWARDHR
C. elegans	VFGGFGITAG	AHRLW <u>SHKSY</u>	KATTPMRIFL	MILNNIALQN	DVIEWARDHR
S.cerevisiae	AVGGVSITAG	YHRLW <u>SHRSY</u>	SAHWPLRLFY	AIFGCASVEG	SAKWGWHSHR
P. angusta	CFGGISITAG	YHRHWA <u>HRAY</u>	DCKLPVKIFF	ALFGASAVEG	SIKMWGHQHR
H. capsulatum	FMTGLGITAG	YHRLWA <u>HCSY</u>	SATLPLKIYL	AAVGGGAVEG	SIRWWARGHR
M. rouxii	FITGLGITAG	YHRMWA <u>SHRAY</u>	RGTDLLRWFM	SFAGAGAVEG	SIYWWSRGHR
C. curvatus	FITGLGITAG	YHRLWA <u>HRSY</u>	NASKPLQYFL	ALCGAGSVQG	SIRWWWSRGHR
C. merolae	FCCGLGITGG	YHRLW <u>SHRSY</u>	EAHWLVQVIL	ACFGAAAFEG	SARYWCLRHR
	201				250
Rat	AHHKFSE <u>THA</u>	DPHNSRRGFF	FSHVGWLLVR	KHPAVKEKGG	KLDMSDLKAE
Mouse	AHHKFSE <u>THA</u>	DPHNSRRGFF	FSHVGWLLVR	KHPAVKEKGG	KLDMSDLKAE
Sheep	AHHKFSE <u>TD</u> A	DPHNSRRGFF	FSHVGWLLVR	KHPAVREKGA	TLDLSDLRAE
Pig	AHHKFSE <u>TD</u> A	DPHNSRRGFF	FSHVGWLLVR	KHPAVKEKGG	LLNMSDLKAE
Human	AHHKFSE <u>THA</u>	DPHNSRRGFF	FSHVGWLLVR	KHPAVKEKGS	TLDLS <u>D</u> LEAE
Hamster	AHHKFSE <u>TYA</u>	DPHNSRRGFF	FSHVGWLLVR	KHPAVKEKGG	KLDMSDLKAE
Drosophila	VHHKYSE <u>TD</u> A	DPHNATRGFF	FSHVGWLLCK	KHPEVKAKGK	GVDLSDLRAD
Moth	MHHKYSE <u>TD</u> A	DPHNATRGFF	FSHVGWLLVR	KHPQIKAKGH	TIDLSDLKSD
C. elegans	CHHKWTD <u>TD</u> A	DPHNTTRGFF	FAHMGWLLVR	KHPQVKEQGA	KLDMSDLLSD
S.cerevisiae	IHHRYTD <u>TLR</u>	DPYDARRGLW	YSHMGWMLLK	PNP...KYKA	RADITDMTDD
P. angusta	VHHRYTD <u>TPR</u>	DPYDAKRGFW	YSHMGWMLLV	PNP...RYKA	RADISDLLDD
H. capsulatum	AHHRYTD <u>TDK</u>	DPYSVRKGLL	YSHIGWMVMK	QNP...KRIG	RTEITDLNED
M. rouxii	AHHRYTD <u>TDK</u>	DPYSAHRGFF	FSHFGWMLVQ	RPK...NRIG	YADVADLKAD
C. curvatus	AHHRYTD <u>TKL</u>	DPYSAHEGFW	HAHMGWMLI.	KPR...GKIG	VADISDLSKN
C. merolae	AHHRYVD <u>SDR</u>	DPYAVEKGFW	YAHLWWMVFK	LPR...QRQG	RVDITDLNAN
	251				300
Rat	KLVMFQRRYY	KPGLLLMCFI	LPTLVPWYCW	GETFLHSLFV	STFLRYTLVL
Mouse	KLVMFQRRYY	KPGLLLMCFI	LPTLVPWYCW	GETFVNSLFV	STFLRYTLVL
Sheep	KLVMFQRRYY	KPGVLLLCFI	LPTLVPWYLW	GESFQNSLFF	ATFLRYAVVL
Pig	KLVMFQRRYY	KPGILLMCFI	LPTIVPWYCW	GEAFQPSLFV	ATFLRYAIVL
Human	KLVMFQRRYY	KPGLLLMCFI	LPTLVPWYFW	GETFQNSVFW	ATFLRYAVVL
Hamster	KLVMFQRRYY	KPAILLMCFI	LPTFVPWYFW	GEAFVNSLCV	STFLRYTLVL
Drosophila	PILMFQKKYY	MILMPIACFI	IPTVVPYAW	GESFMNAFW	ATMFRWCFIL
Moth	PILRFQKKYY	LTLMP LICFI	LPSYIPT.LW	GESAFNAFFV	CSIFRYVYVL
C. elegans	PVLVFQQRHY	FPLVILCCFI	LPTIIPVYFW	KETAFIAFYT	AGTFRYCFTL
S.cerevisiae	WTIRFQHRHY	ILLMLLTAFV	IPTLICGYFF	ND.YMGGLIY	AGFIRVFVIQ
P. angusta	WVVRVQHRHY	LLLMVMAFL	FPAVLTHYLF	ND.FWGGFIY	AGLLRAVVIQ
H. capsulatum	PVVVWQHRNY	LKVVFIMGIV	FPMLVSGLGW	GD.WFGGFIY	AGILRIFFVQ
M. rouxii	HVVAFQHKYY	PYFALGMGFI	FPRTLAVAGLW	GD.FRGGYFY	AGVLRLCFVH
C. curvatus	PVVKWQHNNY	VALLFFMGLA	FPRTLAVAGLW	GD.WWGG LFF	AGAARLVFVH
C. merolae	PILRFQHRYY	LQIAILFSFV	IPLTISTLGW	GD.FWGG L VY	ACLGRMLFVQ

-52-

	301				350
Rat	NATWLVNSAA	HLYGYPYDK	NIQSRNIV	SLGSVGEF	NYHHAFPYDY
Mouse	NATWLVNSAA	HLYGYPYDK	NIQSRNIV	SLGAVGEF	NYHHTFPFDY
Sheep	NATWLVNSAA	HMYPYDK	TINPRENIV	SLGAVGEF	NYHHTFPYDY
Pig	NATWLVNSAA	HLYGYPYDK	TISPRENIV	SLGAVGEF	NYHHTFPYDY
Human	NATWLVNSAA	HLFGYPYDK	NISPRENIV	SLGAVGEF	NYHHSFPYDY
Hamster	NATWLVNSAA	HLYGYPYDK	NIDPRENALV	SLGCLGEF	NYHHAFPYDY
Drosophila	NVTWLVNSAA	HKFGGRPYDK	FINPSENISV	AILAFGEWH	NYHHVFPWDY
Moth	NVTWLVNSAA	HLWGSKPYDK	NINPVETRPV	SLVVLGEF	NYHHTFPWDY
C. elegans	HATWCINSAA	HYFGWKPYDS	SITPVENVFT	TIAAVGEGGH	NFHHTFPQDY
S.cerevisiae	QATFCINSLA	HYIGTQPFDD	RRTPRDNWIT	AIPTFGEYH	NFHHEFPQDY
P. angusta	QATFCVNSLA	HWIGEQPFDD	RRTPRDHVLT	ALPTFGEYH	NFHHEFPQDY
H. capsulatum	QATFCVNSLA	HWLGDPFDD	RNSPRDHIVT	ALVTLGEYH	NFHHEFPQDY
M. rouxii	HATFCVNSLA	HYLGESTFDD	HNTPRDSWVT	ALVTMGEGYH	NFHHTFPQDY
C. curvatus	HSTFCVNSLA	HWLGESTFDD	KHTPKDHFIT	ALVTMGEGYH	NFHHTFPQDY
C. merolae	QSTFCVNSLA	HWGGEQTFSR	RHTSYDSVIT	ALVTLGEYH	NFHHEFPQDY
	351				400
Rat	SASEY.RWHI	NFTTFFIDCM	AALGLAYDRK	KVSKAAVLAR	IKRTGDGSHK
Mouse	SASEY.RWHI	NFTTFFIDCM	AALGLAYDRK	KVSKATVLAR	IKRTGDGSHK
Sheep	SASEY.RWHI	NFTTFFIDCM	AAIGLAYDRK	KVSKAAVLGR	MKRTGEESYK
Pig	SASEY.RWHI	NFTTFFIDCM	AALGLAYDRK	KVSKAAIL--	-----
Human	SASEY.RWHI	NFTTFFIDCM	AALGLAYDRK	KVSKAAILAR	IKRTGDGNYK
Hamster	SASEY.RWHI	NFTTFFIDCM	AALGLAYDRK	KVSKAAVLAR	IKRTGDGSHK
Drosophila	KTAEFGKYSL	NFTTAFIDFF	AKIGWAYDLK	TVSTDIIKKR	VKRTGDGTHA
Moth	KTAEFGKYSL	NFTTAFIDFF	AKIGWAYDLK	TVSTDIIKKR	VKRTGDGTHA
C. elegans	RTSEYS.LKY	NWTRVLIDTA	AALGLVYDRK	TACDEIIGRO	VSNHGCIDQR
S.cerevisiae	RNA.IKQYQY	DPTKVIIYLT	SLVGLAYDLK	KFSQNAIEEA	LIQQEQKKIN
P. angusta	RNA.LKQYQY	DPTKVIIYLT	SKVGLAYNLK	KFSQNAIDQG	ILQQQKKLD
H. capsulatum	RNA.IEWHQY	DPTKWTIWIW	KQLGLAYDLK	QFRANIEIEG	RVQQLQKKID
M. rouxii	RNA.IKQYQY	DPTKWTIWIW	SWFGLAYELK	QFPTNEVTKG	RLFMEEKRIQ
C. curvatus	RNA.IKQYQY	DPTKWTIWIW	AQLGLASHLK	KFPDNEIKKG	QYTMKLMQLQ
C. merolae	RNG.VVWYHW	DPTKWTIWIW	SWAGLAWHLV	RFPRNELVKA	RLQVRQEILD
	401				450
Rat	SS*-----	-----	-----	-----	-----
Mouse	SS-----	-----	-----	-----	-----
Sheep	SG-----	-----	-----	-----	-----
Pig	SG-----	-----	-----	-----	-----
Human	SG-----	-----	-----	-----	-----
Hamster	SG-----	-----	-----	-----	-----
Drosophila	TWGWGDVDQP	KEEIE.DAVI	THKKSE----	-----	-----
Moth	VWGWDDHEVH	QEDKKLAII	NPEKTE----	-----	-----
C. elegans	GKSIM-----	-----	-----	-----	-----
S.cerevisiae	KKKAKINWGP	VLTDLPMDK	QTFLAKS.KE	NKGLVVISGI	VHDSVSGYISE
P. angusta	RMRAKLWGP	QLSELPVWDK	STFFEKA.KE	QKGLVVISGI	VHDCANFLTE
H. capsulatum	QRRAKLDWGI	PLEQLPVIEW	DDYVDQA.KN	GRGLIAIAGV	VHDTVDFIKD
M. rouxii	AQKAKLSYGT	PLKDLPIYTW	EEYQSLVLND	NKKWVLIEGV	LYDVEEFMKE
C. curvatus	EQSEKLEWPK	HSNDLPVISW	EDFQA..ESK	TRALIAVHGF	IHDCCSFIED
C. merolae	EAKKRVWDGK	PIESLPVWTW	KDVQRLAKEE	NRLLVVIEGI	VHDCRTRFKVQ

-53-

	451				500
Rat	-----	-----	-----	-----	-----
Mouse	-----	-----	-----	-----	-----
Sheep	-----	-----	-----	-----	-----
Pig	-----	-----	-----	-----	-----
Human	-----	-----	-----	-----	-----
Hamster	-----	-----	-----	-----	-----
Drosophila	-----	-----	-----	-----	-----
Moth	-----	-----	-----	-----	-----
C. elegans	-----	-----	-----	-----	-----
S. cerevisiae	HPGGETLIKT	ALGKDATKAF	SGGVYRHSNA	AQNVLADMRV	AVIKESKNSA
P. angusta	HPGGQALLKT	SFGKDATMAF	NGGVYAHSNA	AHNLLATMRV	AVIRDGGANG
H. capsulatum	HPGGKAMINS	GIGKDATAMF	NGGVYNHSNA	AHNQLSTMRV	GVIRGGCEVE
M. rouxii	HPGGMKYLST	AVGKDMTTAF	NGGIYNHSNG	TRNLLTSLRV	GVLRNGMQV.
C. curvatus	HPGGAHLIKR	AIGTDSTTAF	FGGVYDHSNA	AHNLLAMMRV	GVLDGGMEVE
C. merolae	HPGGQRILEF	WNVRDATQAF	NGDVYNHTKA	ARNLLAHLRV	AQLKEIYEPE

Protein kinase (specifically cAMP- and cGMP-dependent) phosphorylation sites. There have been a number of studies relative to the specificity of cAMP- and cGMP-dependent protein kinases (Fremisco J.R. et al., J. Biol. Chem. 255:4240-4245, 1980; Glass D.B., Smith S.B., J. Biol. Chem. 258:14797-14803, 1983; Glass D.B. et al., J. Biol. Chem. 261:2987-2993, 1986). Both types of kinases appear to share a preference for the phosphorylation of serine or threonine residues found close to at least two consecutive N-terminal basic residues. It is important to note that there are quite a number of exceptions to this rule. However, the consensus pattern is as follows:

[RK] (2) -x- [ST], where S or T is the phosphorylation site.

Protein kinase C phosphorylation site. In vivo, protein kinase C exhibits a preference for the phosphorylation of serine or threonine residues found close to a C-terminal basic residue (Woodget J.R. et al., Eur. J. Biochem. 161:177-184, 1986; Kishimoto A. et al., J. Biol. Chem. 260:12492-12499, 1985). The presence of additional basic residues at the N- or C-terminus of the target amino acid enhances the Vmax and Km of the

-54-

phosphorylation reaction. The consensus pattern is:

[ST]-x-[RK] where S or T is the phosphorylation site.

Casein kinase II phosphorylation site. Casein kinase II (CK-2) is a protein serine/threonine kinase whose activity is independent of cyclic nucleotides and calcium. CK-2 phosphorylates many different proteins. The substrate specificity (Pinna L.A., Biochim. Biophys. Acta 1054:267-284, 1990) of this enzyme can be summarized as follows: (1) Under comparable conditions Ser is favored over Thr; (2) an acidic residue (either Asp or Glu) must be present three residues from the C-terminal end of the phosphate acceptor site; (3) additional acidic residues in positions +1, +2, +4, and +5 increase the phosphorylation rate (most physiological substrates have at least one acidic residue in these positions); (4) Asp is preferred to Glu as the provider of acidic determinants; and (5) a basic residue at the N-terminus of the acceptor site decreases the phosphorylation rate, while an acidic one will increase it. The consensus pattern is: [ST]-x(2)-[DE] where S or T is the phosphorylation site (note: this pattern is found in most of the known physiological substrates).

If phosphorylation of a specific site by any kinase is found to increase the catalytic activity or stability of the encoded desaturase protein, the phosphorylated serine or threonine residue is changed to encode a negatively charged amino acid (aspartic acid or glutamic acid) in order to permanently optimize the activity or the protein. If phosphorylation of a specific residue is found to decrease the activity or stability of the encoded desaturase, the affected serine or threonine encoding codon is altered to substitute a neutral or a positively charged amino acid that will permanently optimize the activity or stability of the protein.

-55-

EXAMPLE 4**Further Modifications and Improvements of the
Saccharomyces cerevisiae OLE1 Gene for Plant Expression
Using Elements Derived from Native Plant Desaturase Genes**

The activity of the native or modified forms of the *Saccharomyces cerevisiae* OLE1 Δ -9 desaturase gene in plant tissues may be further improved by the substitution or inclusion of elements derived from native plant desaturase genes. Favorable plant gene elements may include sequences that improve the expression of the modified gene at one or more levels, including the following: 1) transcription, 2) pre-mRNA processing, 3) mRNA transport from the nucleus to the cytoplasm, 4) mRNA stability 5) translation, 6) targeting or retention of the protein at the appropriate membrane surface or organelle surface, 7) protein folding and maturation, and 8) stability of the functional desaturase protein.

The inventors have shown that the OLE1 gene can tolerate significant modifications without losing its biological activity. These modifications include deletion of the "coiled coil" region, the addition of 239 amino acids to the N-terminus of OLE1p and truncation of 55 and 60 amino acids from the N-terminal end of the protein. The inventors have also shown that modifications of the 5' and 3'untranslated regions of the OLE1 mRNA can significantly affect its stability. For example, removing a short open reading frame near the 5' "cap" region of the OLE1 mRNA increases its half-life in *Saccharomyces* from 12 minutes to approximately 25 minutes. The existence of elements in the mRNA that affect its stability indicate that other elements might also exist that affect the stability of an mRNA generated by a synthetic gene in another host organism.

Plant desaturase gene elements that enhance the

-56-

function of the modified Δ -9 desaturase gene are identified by a 2-step method. STEP 1 involves isolating a series of DNA sequences from a cDNA that encodes a plant ER lipid biosynthetic enzyme. Those elements are linked, or
5 inserted into regions of a native or "optimized" gene under control of a yeast promoter in a vector suitable for expression in *Saccharomyces cerevisiae*. The resulting vectors are then tested for their ability to produce functional desaturase enzymes in strains of *Saccharomyces*
10 that contain an inactive form of the Δ -9 fatty acid desaturase gene.

In STEP 2, plant desaturase sequences from the above vectors that are found to produce a functional Δ -9 desaturase gene are used to isolate homologous sequences
15 from plant genomic DNA. The isolated genomic sequences are used to construct a synthetic gene that produces an mRNA that encodes the same functional desaturase protein produced by the vector in step 1. In this instance, the genomic sequences encompass the same protein coding
20 elements as those encoded by the homologous cDNA sequence and also include genomic elements that encode the 5' and/or 3' untranslated regions of the plant desaturase mRNA. These combined genomic elements should differ from the cDNA derived sequences used in STEP 1 by containing authentic
25 plant introns, (which may facilitate efficient and correct splicing of the chimeric mRNA in the plant nucleus) and signals that affect the mRNA stability, mRNA transport, and efficient translation of the mRNA in plant tissues. The chimeric plant / synthetic gene containing the genomic
30 sequences is inserted into vectors under the control of plant seed-specific promoters and tested for expression and desaturase function in plants, including *Brassica*, *Arabidopsis*, maize and soybeans.

-57-

The following specific examples further illustrate these methods employing the *Arabidopsis FAD2* gene, which encodes an ER $\Delta 12$ -desaturase, as a source of plant desaturase DNA sequences. In the preferred embodiment, the source of the plant desaturase DNA would be the *FAD2* homolog, or a related ER lipid biosynthetic gene, that is derived from the same plant species that is intended to be modified by the resulting vector for commercial use.

10

A. Substitution of the N-terminal *OLE1* protein coding sequences and with N-terminal sequences from the derived from the *Arabidopsis FAD2* gene.

1) A cDNA containing the *FAD2*, Δ -12 desaturase, mRNA coding sequence is isolated by reverse transcriptase - polymerase chain reaction (RT-PCR) of isolated mRNAs derived from *Arabidopsis* tissue or by direct DNA synthesis using the protein and DNA sequences set forth in SEQ ID NO:4 and SEQ ID NO:5 (open reading frame starts at +93).

2) The inventors have shown that substitution of transmembrane sequences of the *OLE1* gene with transmembrane sequences from the *Saccharomyces FAH2* gene abolishes Δ -9 desaturase activity. *FAH2* encodes a sphingolipid fatty acid hydroxylase, which is an ER membrane protein.

TMPredict analysis of the *Arabidopsis FAD2* sequence indicates that the first transmembrane region of its encoded protein begins at residue +52 and a similar analysis of the *OLE1* sequence indicates that its first transmembrane sequence begins at residue +113. Because the inclusion of potential membrane spanning elements from the plant desaturase could produce significant changes in the desaturase core enzyme structure that affect activity, only sequences encoding residues +1 to +52 of *FAD2* are tested

-58-

for functional linkages or substitutions in the 113 residue N-terminal region of *OLE1*.

A series of PCR oligonucleotide primers are synthesized that include a 5' primer that complements sequences including +1 start codon of the *FAD2* gene and 3' primers that complement sequences ending, for example, at residues +20, +35 and + 52 of the *FAD2* gene. These are used to amplify a series of fragments of different lengths from the *FAD2* cDNA that extend from the +1 codon through codon +52. A second PCR amplification is performed using a 5' primer that is complementary to sequences that include the 5' end of the *FAD2* mRNA and the 3' primer that includes codon +52. That amplification is done using *Arabidopsis* genomic DNA as a template. The amplified fragment from that reaction is cloned into a bacterial vector and subjected to DNA sequencing to detect the presence of introns within the genomic sequence. The cloned genomic fragment is also used to construct vectors for plant expression as indicated in STEP 2 of the method.

The amplified cDNA fragments is inserted into yeast expression vectors that contain the native *OLE1* mRNA coding sequence under the control of the *Saccharomyces* galactose inducible, *GAL1* promoter. Insertion of the plant DNA fragments can be done in several ways: 1) A fragment is inserted upstream of the *OLE1* protein coding sequences so that its protein coding element is fused in frame to the +1 codon of the *OLE1* encoded protein, 2) the codons on the plant fragment could replace the equivalent *OLE1* residues starting from the +1 ATG codon (e.g. a plant DNA fragment containing codons +1 -> +52 replaces *OLE1* codons +1 -> +52) and 3) the full length fragment containing codons +1 -> + 52 of the plant gene is fused in frame to codon +114 of the *OLE1* gene, replacing the *OLE1* residues +1 -> +113 with

-59-

plant desaturase residues +1 -> +52.

The resulting plasmids are transformed into a haploid *ole1Δ::LEU2* strain of *Saccharomyces*. That strain contains a null, disrupted form of the *OLE1* gene and
5 therefore has a growth requirement for unsaturated fatty acids. The transformed *Saccharomyces* strains are grown on fatty acid depleted galactose medium to test for the ability of the induced chimeric gene to support growth of the strain without fatty acids. Transformed strains that
10 grow on the fatty acid deficient medium are further analyzed to assess the effects of the plant sequences on desaturase function. This is done by Western blot analysis, to measure levels of the resulting desaturase protein and by fatty acid analysis of total cellular
15 lipids, to assess the relative activity of the desaturase enzyme by comparison of the ratio of saturated to unsaturated fatty acids.

3) Using information derived from the above tests, a chimeric desaturase gene is constructed using the
20 amplified genomic DNA from the *FAD2* gene. Construction, testing, and analysis these vectors is guided by the principle that the most desirable vector is one that maximizes the use of the plant gene sequences and minimizes the use of the *Saccharomyces* Δ-9 desaturase gene sequences
25 while retaining optimal desaturase function. Plant DNA fragments derived from the genomic DNA amplification that extend from the 5' end of the mRNA sequence to the longest sequence that produces optimal desaturase function in yeast are inserted into a vector containing the native Δ-9
30 desaturase gene (or one of its modified forms produced by the methods described above). The fragment is inserted into the vector so that the 3' end of its protein coding sequence produces an mRNA that generates a protein sequence

-60-

identical to its counterpart derived from the *FAD2* cDNA sequences. The resulting chimeric desaturase gene, which now encodes an mRNA that includes the *FAD2* 5' untranslated region in addition to the modified protein coding sequences, is placed into a plant expression vector under the control of a suitable plant promoter and plant termination/ polyadenylation sequences.

4) The resulting vectors containing the plant/yeast chimeric desaturase sequences are transformed into plants for testing and analysis of desaturase function. Suitable test plants include *Arabidopsis thaliana*, and *Brassica napis*. A method for transformation and analysis of desaturase gene expression in *Arabidopsis* is provided above. A method for transformation and analysis of yeast desaturase expression in *Brassica napis* is described in U.S. Patent No. 5,777,201 to Poutre et al. (incorporated by reference herein).

B. Insertion or substitution of *Arabidopsis FAD2* C-terminal protein coding sequences and 3' mRNA untranslated region sequences into native and modified forms of the *OLE1* gene.

The inventors have previously shown that proteins encoded by the *Saccharomyces ELO2* and *ELO3* genes contain a series of charged residues in their C-terminal region. These proteins are located on the ER surface and function in the biosynthesis of very long chain fatty acids as described in Oh et al. (J. Biol. Chem. 272: 17376-17384, 1997) (incorporated by reference herein). They further showed that deletion of the region containing the charged residues causes the proteins to be mislocalized from their normal cellular locations in the endoplasmic reticulum, resulting in reduced function. Similar clusters

-61-

of charged residues occurs in the C-terminal region of the *OLE1* gene that are apparently associated with ER retention or localization. These residues do not appear to be a part of the functional cytochrome *b₅* domain. A detailed

5 comparison of the C-terminal *OLE1* and the Arabidopsis *FAD2* sequences show that the plant desaturase has similar, but not identical, clusters of charged residues to those in the *OLE1* gene. These sequences are shown below:

10 SEQ ID Nos: 6 and 7:

Comparison of the charged carboxyl terminal amino acids of Ole1p (SEQ ID NO:7) and the Arabidopsis Fad2p desaturase (SEQ ID NO:6) (The region of the *OLE1* gene shown does not appear to be a functional part of its cytochrome *b₅*,
15 domain).

A.thaliana FAD2 +- + - - -+- -++ +
WYVAMYREAK ECIYVEPDRE GDKKGVYWYN NKL*

20 *S.cerevisiae OLE1* +- + + ++ - - +
MRVAVIKESK NSAIRMASKR GEIYETGKFF *

Methods similar to those shown in Section A can be used to
25 identify Arabidopsis *FAD2* sequences that can replace the *OLE1* C-terminal sequences to optimize gene expression, membrane targeting and ER retention of the chimeric enzyme.

1) A series of oligonucleotide primers for PCR
amplification are synthesized for isolation of elements in
30 the C-terminal region of the *FAD2* gene. A *FAD2* DNA fragment encompassing that region is generated by PCR amplification of the cDNA clone. Alternatively, given the smaller size of the fragment it or modified forms of the plant fragment may be generated directly by DNA synthesis.

-62-

A fragment containing that region and its flanking 3' untranslated region also is generated by PCR amplification of *Arabidopsis* genomic DNA as described above. That fragment is cloned into an appropriate vector and sequenced. as also described.

2) Vectors are constructed that contain the plant DNA fragments linked to or substituted into the *OLE1* C-terminal coding region as described in Section A. In this instance, the plant DNA fragments are linked in frame to the carboxyl terminal residues of the *OLE1* protein coding region.

3) The resulting vectors are transformed into the *Saccharomyces ole1Δ* strain and tested for desaturase function as described in Section A.

4) Using information derived from the above tests, chimeric desaturase genes containing the C-terminal plant sequences that produce functional desaturases are constructed using the amplified genomic DNA from the *FAD2* gene, according to the principles outlined in Section A. The resulting sequences are employed to construct vectors that will express the chimeric plant/yeast gene under control of plant promoter and plant termination/polyadenylation sequences. Those vectors are transformed into plants for testing and analysis of desaturase function as described above.

The present invention is not limited to the embodiments described and exemplified above, but is capable of variation and modification without departure from the scope of the appended claims.

-63-

We claim:

1. A synthetic fatty acid desaturase gene for
5 expression in a multicellular plant, the gene comprising a
desaturase domain and a cyt b_5 domain, wherein the gene is
customized for expression in a plant cytoplasm.
2. The synthetic gene of claim 1, customized
10 from a naturally occurring gene encoding a cytosolic Δ -9
desaturase.
3. The synthetic gene of claim 2, customized
from a naturally occurring gene from *Saccharomyces*
15 *cerevisiae*.
4. The synthetic gene of claim 3, customized
from a naturally occurring gene from *Saccharomyces*
cerevisiae that encodes SEQ ID NO:2.
20
5. The synthetic gene of claim 4, customized
from a naturally occurring gene from *Saccharomyces*
cerevisiae comprising SEQ ID NO:1.
- 25 6. The synthetic gene of claim 3, comprising SEQ
ID NO:3.
7. The synthetic gene of claim 1, which further
comprises an expression regulatory sequence from a plant
30 gene encoding an ER biosynthetic pathway enzyme.
8. The synthetic gene of claim 1, customized for
expression in a monocotyledonous plant.

-64-

9. The synthetic gene of claim 1, customized for expression in a dicotyledonous plant.

10. The synthetic gene of claim 1, customized
5 for expression in a plant genus selected from the group consisting of *Arabidopsis*, *Brassica*, *Phaseolus*, *Oryza*, *Olea*, *Elaeis* (Oil Palm) and *Zea*.

11. The synthetic gene of claim 1, customized
10 from a naturally occurring gene comprising both a desaturase domain and a cyt b₅ domain.

12. The synthetic gene of claim 1, wherein the gene is a chimeric gene comprising a desaturase domain and
15 a heterologous cyt b₅ domain.

13. The synthetic gene of claim 1, customized from a naturally occurring gene such that the synthetic gene and the naturally occurring gene encode an identical
20 amino acid sequence.

14. The synthetic gene of claim 13, wherein the synthetic gene and the naturally occurring gene encode SEQ ID NO:2.
25

15. The synthetic gene of claim 1, customized from a naturally occurring gene such that the synthetic gene and the naturally occurring gene encode a similar amino acid sequence.
30

16. The synthetic gene of claim 1, customized from a naturally occurring gene such that the synthetic gene and the naturally occurring gene encode a similar

-65-

amino acid sequence, and the synthetic gene possesses improved stability or catalytic activity as compared with the naturally occurring gene.

5 17. A method for constructing a customized bifunctional desaturase/cyt b_5 encoding gene for expression in the cytosol of a multicellular plant, comprising the steps of:

 (a) providing a DNA molecule comprising a
10 desaturase-encoding moiety operably linked to a cyt b_5 -
encoding moiety, said DNA molecule producing the
bifunctional polypeptide in a non-customized form;

 (b) back-translating the polypeptide
sequence using preferred codons for expression in a
15 multicellular plant, thereby producing a back-translated
nucleotide sequence;

 (c) analyzing the back-translated nucleotide
sequence for features that could diminish or prevent
expression in the plant cytoplasm;

20 (d) modifying the analyzed sequence to
correct or remove the features that could diminish or
prevent expression in the plant cytoplasm; and

 (e) optionally, introducing pre-determined
cloning features into the sequence in a manner that does
25 not materially affect the codon usage or final polypeptide
sequence, thereby producing the customized bifunctional
desaturase/cyt b_5 encoding gene for expression in the
cytosol of a multicellular plant.

30 18. The method of claim 17, wherein the features
that could diminish or prevent expression in the plant
cytoplasm include one or more features selected from the
group consisting of: putative intron splice sites, plant

-66-

polyadenylation signals, RNA polymerase II termination sequences, and hairpin consensus sequences.

19. The method of claim 17, which further
5 comprises the step of:

(f) testing the customized bifunctional desaturase/cyt b₅ encoding gene for desaturase function in fatty acid deficient strains of a microorganism prior introducing the gene into vectors for expression in plants.

10

20. The method of claim 19, wherein the microorganism is *Saccharomyces cerevisiae*.

21. The method of claim 17, which further
15 comprises incorporating into the customized gene one or more genomic segments from plant desaturase or other ER lipid biosynthetic genes, which comprise beneficial elements to further optimize expression of the genes in plants, comprising the steps of:

20 a) selecting a cDNA sequence that potentially comprises one or more of the beneficial elements;

b) creating a yeast vector expressing a desaturase gene modified to contain one or more of the
25 beneficial elements;

c) testing the vector in a yeast expression system;

d) isolating regions from genomic DNA that are homologous to the beneficial elements from the cDNA;
30 and

e) operably linking the genomic DNA regions to the customized bifunctional desaturase/cyt b₅ encoding gene to produce the further customized gene.

1/2




FIGURE 1

2/2




FIGURE 2

SEQUENCE LISTING

<110> Martin, Charles E.
Mitchell, Andrew

<120> Synthetic Fatty Acid Desaturase Gene for
Expression in Plants

<130> 97-0081 PCT

<150> US 60/097,586

<151> 1998-08-24

<160> 7

<170> FastSEQ for Windows Version 3.0

<210> 1

<211> 1555

<212> DNA

<213> *Saccharomyces cerevisiae*

<400> 1

tacaacaaag	atgccaaactt	ctggaactac	tattgaattg	attgacgacc	aatttccaaa	60
ggatgactct	gccagcagtg	gcattgtcga	cgaagtcgac	ttaacggaag	ctaataat	120
ggctactggt	ttgaataaga	aagcaccaag	aattgtcaac	ggttttgggt	ctttaatggg	180
ctccaaggaa	atggtttccg	tggaattcga	caagaaggga	aacgaaaaga	agtccaat	240
ggatcgctctg	ctagaaaagg	acaaccaaga	aaaagaagaa	gctaaaacta	aaattcacat	300
ctccgaacaa	ccatggactt	tgaataactg	gcaccaacat	ttgaactggg	tgaacatggg	360
tcttgtttgt	ggtatgcaa	tgattgggtg	gtacttcgct	ctctctggta	aagtacctt	420
gcatttaaac	gttttccttt	tctccgtttt	ctactacgct	gtcgggtggg	tttctattac	480
tgccgggtac	catagattat	ggtctcacag	atcttactcc	gctcactggc	cattgagatt	540
attctacgct	atcttcgggt	gtgcttccgt	tgaagggtcc	gctaaatggg	ggggccactc	600
tcacagaatt	caccatcggt	acactgatac	cttgagagat	ccttatgacg	ctcgtagagg	660
tctatgggtac	tcccacatgg	gatggatgct	tttgaagcca	aatccaaaat	acaaggctag	720
agctgatatt	accgatatga	ctgatgattg	gaccattaga	ttccaacaca	gacactacat	780
cttggtgatg	ttattaaccg	ctttcgtcat	tccaactctt	atctgtgggt	actttttcaa	840
cgactatatg	ggtgggttga	tctatgccgg	ttttattcgt	gtctttgtca	ttcaacaagc	900
taccttttgc	attaactcca	tggtcatta	catcggtacc	caaccattcg	atgacagaag	960
aaccctctgt	gacaactgga	ttactgccat	tgttactttc	ggtgaagggt	accataactt	1020
ccaccacgaa	ttcccaactg	attacagaaa	cgctattaag	tggtaccaat	acgacccaac	1080
taaggttatc	atctatattga	cttctttagt	tggtctagca	tacgacttga	agaaattctc	1140
tcaaaatgct	attgaagaag	ccttgattca	acaagaacaa	aagaagatca	ataaaaagaa	1200
ggctaagatt	aactggggtc	cagttttgac	tgatttgcca	atgtgggaca	aacaaaacct	1260
cttggttaag	tctaaggaaa	acaagggttt	ggttatcatt	tctgggtattg	ttcacgacgt	1320
atctgggttat	atctctgaac	atccaggtgg	tgaaacttta	attaaaactg	cattaggttaa	1380
ggacgctacc	aaggctttca	gtgggtgggt	ctaccgtcac	tcaaatgccg	ctcaaatgtg	1440
cttggctgat	atgagagtgg	ctgttatcaa	ggaaagtaag	aactctgcta	ttagaatggc	1500
tagtaagaga	ggtgaaatct	acgaaactgg	taagttcttt	taagcatcac	attac	1555

<210> 2

<211> 510

<212> PRT

<213> *Saccharomyces cerevisiae*

<400> 2

Met Pro Thr Ser Gly Thr Thr Ile Glu Leu Ile Asp Asp Gln Phe Pro
 1 5 10 15
 Lys Asp Asp Ser Ala Ser Ser Gly Ile Val Asp Glu Val Asp Leu Thr
 20 25 30
 Glu Ala Asn Ile Leu Ala Thr Gly Leu Asn Lys Lys Ala Pro Arg Ile
 35 40 45
 Val Asn Gly Phe Gly Ser Leu Met Gly Ser Lys Glu Met Val Ser Val
 50 55 60
 Glu Phe Asp Lys Lys Gly Asn Glu Lys Lys Ser Asn Leu Asp Arg Leu
 65 70 75 80
 Leu Glu Lys Asp Asn Gln Glu Lys Glu Glu Ala Lys Thr Lys Ile His
 85 90 95
 Ile Ser Glu Gln Pro Trp Thr Leu Asn Asn Trp His Gln His Leu Asn
 100 105 110
 Trp Leu Asn Met Val Leu Val Cys Gly Met Pro Met Ile Gly Trp Tyr
 115 120 125
 Phe Ala Leu Ser Gly Lys Val Pro Leu His Leu Asn Val Phe Leu Phe
 130 135 140
 Ser Val Phe Tyr Tyr Ala Val Gly Gly Val Ser Ile Thr Ala Gly Tyr
 145 150 155 160
 His Arg Leu Trp Ser His Arg Ser Tyr Ser Ala His Trp Pro Leu Arg
 165 170 175
 Leu Phe Tyr Ala Ile Phe Gly Cys Ala Ser Val Glu Gly Ser Ala Lys
 180 185 190
 Trp Trp Gly His Ser His Arg Ile His His Arg Tyr Thr Asp Thr Leu
 195 200 205
 Arg Asp Pro Tyr Asp Ala Arg Arg Gly Leu Trp Tyr Ser His Met Gly
 210 215 220
 Trp Met Leu Leu Lys Pro Asn Pro Lys Tyr Lys Ala Arg Ala Asp Ile
 225 230 235 240
 Thr Asp Met Thr Asp Asp Trp Thr Ile Arg Phe Gln His Arg His Tyr
 245 250 255
 Ile Leu Leu Met Leu Leu Thr Ala Phe Val Ile Pro Thr Leu Ile Cys
 260 265 270
 Gly Tyr Phe Phe Asn Asp Tyr Met Gly Gly Leu Ile Tyr Ala Gly Phe
 275 280 285
 Ile Arg Val Phe Val Ile Gln Gln Ala Thr Phe Cys Ile Asn Ser Leu
 290 295 300
 Ala His Tyr Ile Gly Thr Gln Pro Phe Asp Asp Arg Arg Thr Pro Arg
 305 310 315 320
 Asp Asn Trp Ile Thr Ala Ile Val Thr Phe Gly Glu Gly Tyr His Asn
 325 330 335
 Phe His His Glu Phe Pro Thr Asp Tyr Arg Asn Ala Ile Lys Trp Tyr
 340 345 350
 Gln Tyr Asp Pro Thr Lys Val Ile Tyr Leu Thr Ser Leu Val Gly
 355 360 365
 Leu Ala Tyr Asp Leu Lys Lys Phe Ser Gln Asn Ala Ile Glu Glu Ala
 370 375 380
 Leu Ile Gln Gln Glu Gln Lys Lys Ile Asn Lys Lys Lys Ala Lys Ile
 385 390 395 400
 Asn Trp Gly Pro Val Leu Thr Asp Leu Pro Met Trp Asp Lys Gln Thr
 405 410 415
 Phe Leu Ala Lys Ser Lys Glu Asn Lys Gly Leu Val Ile Ile Ser Gly
 420 425 430
 Ile Val His Asp Val Ser Gly Tyr Ile Ser Glu His Pro Gly Gly Glu
 435 440 445
 Thr Leu Ile Lys Thr Ala Leu Gly Lys Asp Ala Thr Lys Ala Phe Ser
 450 455 460
 Gly Gly Val Tyr Arg His Ser Asn Ala Ala Gln Asn Val Leu Ala Asp

3

465					470					475				480
Met	Arg	Val	Ala	Val	Ile	Lys	Glu	Ser	Lys	Asn	Ser	Ala	Ile	Arg
				485					490					495
Ala	Ser	Lys	Arg	Gly	Glu	Ile	Tyr	Glu	Thr	Gly	Lys	Phe	Phe	
			500					505					510	

```
<210> 3
<211> 1555
<212> DNA
<213> Artificial Sequence
```

<220>
<223> synthetic yeast delta-9 desaturase gene modified
for expression in plants

<400> 3							
ggatccaaca	atgcctactt	ctggaactac	tatcgagctt	atcgatgatc	aattccctaa		60
ggatgattct	gcttcttctg	gaatcgttga	tgaggttgat	cttactgagg	ctaactcctt		120
tgctactgga	cttaacaaga	agggctctag	aatcggttaac	ggattcggat	ctcttatggg		180
atctaaggag	atggtttctg	ttaggttctga	taagaaggga	aacgagaaga	agtctaacct		240
tgatagactt	cttgagaagg	ataaccaaga	gaaggaggag	gctaagacta	agatccatat		300
ctctgagcaa	ccttggactc	tcaacaactg	gcataacat	ctcaactggc	tcaacatggt		360
gctcgtctgt	ggaatgccta	tgatcggatg	gtacttgcct	ctctctggaa	aagtgcctct		420
ccatctcaac	gttttctctt	tctctgtctt	ctactacgtt	gttggaggag	tgtctatcac		480
tgtcggatac	catagactct	ggtctcatag	atcttactct	gctcatggc	ctcttagact		540
cttctacgct	atctttggat	gtgcttctgt	tgagggatct	gctaagtggt	ggggacattc		600
tcatagaatc	catcatagat	acactgatac	tcttagagat	ccttacgatg	ctagaagagg		660
actttggtac	tctcatatgg	gatggatgct	tcttaagcct	aaccctaagt	acaaggctag		720
agctgatatc	actgatatga	ctgatgattg	gactatcaga	ttccaacata	gacatttatc		780
cttgctcatg	ctccttactg	ctttctgtat	ccctactctc	atctgtggat	acttctcaa		840
cgattacatg	ggaggactca	tctacgctgg	attcatcaga	gtgttctga	tccaacaagc		900
ctatttctgt	atcaactcta	tggctcata	catcggaa	caacctttcg	atgatagaag		960
aactcctaga	gataactgga	tcaactgctat	cgttactttc	ggagagggat	accataactt		1020
ccatcatgag	ttccctactg	attatagaaa	cgctatcaag	tggtaccaat	acgactctac		1080
taaagtgatc	atctacttga	cttctctcgt	gggacttgct	tacgattctca	agaagtcttc		1140
tcaaaacgct	atcgaggagg	ctcttatcca	acaagagcaa	aagaagatca	acaagaagaa		1200
ggctaagatt	aattggggag	ctgttctttac	tgatcttctt	atgtgggata	agcaaacttt		1260
ccttgctaag	tctaaggaga	acaagggact	tgttatcatc	tctggaatcg	ttcatgatgt		1320
ttctggatac	atctctgagc	atcctggaag	agagacttta	attaagactg	ctcttggaaa		1380
ggatgctact	aaggctttct	ctggaggagt	ttacagacat	cttaacgctg	tcaaaacgct		1440
gcttgctgat	atgagagttg	ctgttatcaa	ggagttctaa	aactctgcta	ctcagaatggc		1500
ttctaaqaga	qgagagactc	acgagactgg	aaagtctctc	tgatctagag	gatcc		1555

```
<210> 4
<211> 383
<212> PRT
<213> Arabidopsis thaliana
```

<400> 4															
Met	Gly	Ala	Gly	Gly	Arg	Met	Pro	Val	Pro	Thr	Ser	Ser	Lys	Lys	Ser
1				5					10					15	
Glu	Thr	Asp	Thr	Thr	Lys	Arg	Val	Pro	Cys	Glu	Lys	Pro	Pro	Phe	Ser
			20					25					30		
Val	Gly	Asp	Leu	Lys	Lys	Ala	Ile	Pro	Pro	His	Cys	Phe	Lys	Arg	Ser
		35					40					45			
Ile	Pro	Arg	Ser	Phe	Ser	Tyr	Leu	Ile	Ser	Asp	Ile	Ile	Ile	Ala	Ser
	50					55				60					
Cys	Phe	Tyr	Tyr	Val	Ala	Thr	Asn	Tyr	Phe	Ser	Leu	Leu	Pro	Gln	Pro
65					70					75					80

Leu Ser Tyr Leu Ala Trp Pro Leu Tyr Trp Ala Cys Gln Gly Cys Val
 85 90 95
 Leu Thr Gly Ile Trp Val Ile Ala His Glu Cys Gly His His Ala Phe
 100 105 110
 Ser Asp Tyr Gln Trp Leu Asp Asp Thr Val Gly Leu Ile Phe His Ser
 115 120 125
 Phe Leu Leu Val Pro Tyr Phe Ser Trp Lys Tyr Ser His Arg Arg His
 130 135 140
 His Ser Asn Thr Gly Ser Leu Glu Arg Asp Glu Val Phe Val Pro Lys
 145 150 155 160
 Gln Lys Ser Ala Ile Lys Trp Tyr Gly Lys Tyr Leu Asn Asn Pro Leu
 165 170 175
 Gly Arg Ile Met Met Leu Thr Val Gln Phe Val Leu Gly Trp Pro Leu
 180 185 190
 Tyr Leu Ala Phe Asn Val Ser Gly Arg Pro Tyr Asp Gly Phe Ala Cys
 195 200 205
 His Phe Phe Pro Asn Ala Pro Ile Tyr Asn Asp Arg Glu Arg Leu Gln
 210 215 220
 Ile Tyr Leu Ser Asp Ala Gly Ile Leu Ala Val Cys Phe Gly Leu Tyr
 225 230 235 240
 Arg Tyr Ala Ala Ala Gln Gly Met Ala Ser Met Ile Cys Leu Tyr Gly
 245 250 255
 Val Pro Leu Leu Ile Val Asn Ala Phe Leu Val Leu Ile Thr Tyr Leu
 260 265 270
 Gln His Thr His Pro Ser Leu Pro His Tyr Asp Ser Ser Glu Trp Asp
 275 280 285
 Trp Leu Arg Gly Ala Leu Ala Thr Val Asp Arg Asp Tyr Gly Ile Leu
 290 295 300
 Asn Lys Val Phe His Asn Ile Thr Asp Thr His Val Ala His His Leu
 305 310 315 320
 Phe Ser Thr Met Pro His Tyr Asn Ala Met Glu Ala Thr Lys Ala Ile
 325 330 335
 Lys Pro Ile Leu Gly Asp Tyr Tyr Gln Phe Asp Gly Thr Pro Trp Tyr
 340 345 350
 Val Ala Met Tyr Arg Glu Ala Lys Glu Cys Ile Tyr Val Glu Pro Asp
 355 360 365
 Arg Glu Gly Asp Lys Lys Gly Val Tyr Trp Tyr Asn Asn Lys Leu
 370 375 380

<210> 5

<211> 1372

<212> DNA

<213> Arabidopsis thaliana

<400> 5

agagagagag attctgcgga ggagcttctt cttcgtaggg tgttcacgtt tattaacggt	60
atcgccccta cgtcagctcc atctccagaa acatgggtgc aggtggaaga atgccggttc	120
ctacttcttc caagaaatcg gaaaccgaca ccacaaagcg tgtgccgtgc gagaaaccgc	180
ctttctcggt gggagatctg aagaaagcaa tcccgccgca ttgtttcaaa cgctcaatcc	240
ctcgctcttt ctectacctt atcagtgcac tcattatagc ctcagtcttc tactacgtcg	300
ccaccaatta cttctctctc ctccctcagc ctctctctta cttggcttgg ccactctatt	360
gggcctgtca aggctgtgtc ctaactggta tctgggtcat agcccacgaa tgcggtcacc	420
acgcatttcag cgactaccaa tggctggatg acacagttgg tcttatcttc cattccttcc	480
tcctcgtecc ttacttctcc tggaggtata gtcacgccc tcaccattcc aacactggat	540
ccctcgaaaag agatgaagta tttgtcccaa agcagaaatc agcaatcaag tggtagcgga	600
aatacctcaa caacctctt ggacgcacat tgatgttaac cgtccagttt gtcctcgggt	660
ggcccttgta cttagccttt aacgtctctg gcagaccgta tgacgggttc gcttgccatt	720
tttcccccac cgctcccatc tacaatgacc gagaacgcct ccagatatat ctctctgatg	780
cgggtattct agcctctctt tttggtcttt accgttacgc tgctgcacaa gggatggcct	840

5

```

cgatgatctg cctctacgga gtaccgcttc tgatagtga tgcgttcctc gtcttgatca 900
cttacttgca gcacactcat ccctcggtgc ctcactacga ttcacagag tgggactggc 960
tcaggggagc tttggctacc gtagacagag actacggaat cttgaacaag gtgttcaca 1020
acattacaga cacacacgtg gtcctacacc tggtctcgac aatgccgcat tataacgcaa 1080
tggaagctac aaaggcgata aagccaattc tgggagacta ttaccagttc gatggaacac 1140
cgtggtatgt agcgatgtat agggaggcaa aggagtgtat ctatgtagaa ccggacaggg 1200
aagggtgaaa gaaagggtgtg tactggtaca acaataagtt atgagcatga tggggaagaa 1260
attgtcgacc tttctcttgt ctgtttgtct tttgttaaag aagctatgct tcgttttaat 1320
aatcttattg tccattttgt tgtgttatga cattttggct gctcattatg tt 1372

```

<210> 6

<211> 33

<212> PRT

<213> *Arabidopsis thaliana*

<400> 7

```

Trp Tyr Val Ala Met Tyr Arg Glu Ala Lys Glu Cys Ile Tyr Val Glu
1           5           10          15
Pro Asp Arg Glu Gly Asp Lys Lys Gly Val Tyr Trp Tyr Asn Asn Lys
20          25          30
Leu

```

<210> 7

<211> 30

<212> PRT

<213> *Saccharomyces cerevisiae*

<400> 7

```

Met Arg Val Ala Val Ile Lys Glu Ser Lys Asn Ser Ala Ile Arg Met
1           5           10          15
Ala Ser Lys Arg Gly Glu Ile Tyr Glu Thr Gly Lys Phe Phe
20          25          30

```

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US99/19443

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) :C07H 21/02; C12N 15/82; A01H 5/00

US CL :536/23.1; 800/281, 298; 435/69.1, 419, 468

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 536/23.1; 800/281, 298; 435/69.1, 419, 468

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

WEST/DERWENT, GENBANK, EMBL

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 5,777,201 A (POUTRE et al) 07 July 1998, especially col.14-18.	1-5,9-11,13-15
-		-----
Y		6-8,12,16-21
Y	US 5,057,419 A (MARTIN et al) 15 October 1991, col. 7-11.	1-21
Y	STUKEY et al. The OLE1 Gene of Saccharomyces cerevisiae Encodes the delta 9 Fatty Acid Desaturase and Can Be Functionally Replaced by the Rat Stearoyl-CoA Desaturase. Gene. The Journal of Biological Chemistry. 25 November 1990, Vol. 265, No. 33, pages 20144-20149, especially page 20146.	1-21

☒ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
B earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*A* document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means	
P document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

12 NOVEMBER 1999

Date of mailing of the international search report

06 DEC 1999

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

D. Lawrence Fox
ELIZABETH F. MCELWAIN

Telephone No. 308-0196

Form PCT/ISA/210 (second sheet)(July 1992)*

INTERNATIONAL SEARCH REPORT

Int. .ational application No.
PCT/US99/19443

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X - Y	POLASHOCK et al. Expression of the Yeast delta-9 Fatty Acid Desaturase in Nicotiana tabacum. Plant Physiology. 1992, Vol. 100, pages 894-901, especially page 895.	1-5,9-11,13-15 ----- 6-8,12,16-21

Form PCT/ISA/210 (continuation of second sheet)(July 1992)*